

# Bioactive Compounds and Health Benefits of *Artemisia* Species

Natural Product Communications  
 July 2019: 1–17  
 © The Author(s) 2019  
 Article reuse guidelines:  
[sagepub.com/journals-permissions](http://sagepub.com/journals-permissions)  
 DOI: 10.1177/1934578X19850354  
[journals.sagepub.com/home/npx](http://journals.sagepub.com/home/npx)



Manisha Nigam<sup>1</sup>, Maria Atanassova<sup>2</sup>, Abhay P. Mishra<sup>3</sup>,  
 Raffaele Pezzani<sup>4,5</sup>, Hari Prasad Devkota<sup>6</sup>, Sergey Plygun<sup>7,8,9</sup>, Bahare Salehi<sup>10</sup>,  
 William N. Setzer<sup>11,12</sup>, and Javad Sharifi-Rad<sup>13,14</sup>

## Abstract

*Artemisia* L. is a genus of small herbs and shrubs found in northern temperate regions. It belongs to the important family Asteraceae, one of the most numerous plant groupings, which comprises about 1000 genera and over 20000 species. *Artemisia* has a broad spectrum of bioactivity, owing to the presence of several active ingredients or secondary metabolites, which work through various modes of action. It has widespread pharmacological activities and has been used as traditional medicine since ancient times as an anthelmintic, anti-spasmodic, antirheumatic, and antibacterial agent and for the treatment of malaria, hepatitis, cancer, inflammation, and menstrual-related disorders. This review comprises the updated information about the ethnomedical uses and health benefits of various *Artemisia* spp. and general information about bioactive compounds and free radicals.

## Keywords

*Artemisia*, bioactive compounds, artemisinin, essential oils

Received: November 11th, 2018; Revised: February 15th, 2019;  
 Accepted: February 19th, 2019.

*Artemisia* L. is included in the tribe Anthemideae and comprises over 500 species, which are mainly found in Asia, Europe, and North America.<sup>1–3</sup> A large number of members of the Anthemideae tribe are important as flowers and ornamental crops, as well as medicinal and aromatic plants, many of which produce essential oils used in folk and modern medicine, and in the cosmetics and pharmaceutical industry.<sup>2,4</sup> The genus *Artemisia* comprises a variable number of species found throughout the northern half of the world. The genus may be divided into sections *Artemisia* and *Dracunculus*.<sup>5–7</sup> *Artemisia* comprises over 400 species, many of which have an aromatic, bitter taste. Some say that it is named after the Greek Artemis, who was the goddess of the hunt, of forests, and of childbirth.<sup>5,8,9</sup>

The large genus *Artemisia* from the tribe Anthemideae comprises important medicinal plants, which are currently the subject of phytochemical attention because of their biological and chemical diversity and essential oil production.<sup>2</sup> *Artemisia* generally has a broad spectrum of bioactivity, owing to the presence of several active ingredients or secondary metabolites, which work through various modes of action. Secondary metabolism in a plant not only plays a role in its survival by producing attractants for pollinators, but also acts as a chemical defense against herbivory and disease.<sup>10,11</sup> *Artemisia* essential oils, mostly distilled from the aromatic plants, contain a variety

<sup>1</sup> Department of Biochemistry, H. N. B. Garhwal University, Srinagar, India

<sup>2</sup> Scientific Consulting, Chemical Engineering, UCTM, Sofia, Bulgaria

<sup>3</sup> Department of Pharmaceutical Chemistry, H. N. B. Garhwal University, Srinagar, India

<sup>4</sup> OU Endocrinology, Department of Medicine (DIMED), University of Padova, Italy

<sup>5</sup> AIROB, Associazione Italiana per la Ricerca Oncologica di Base, Padova, Italy

<sup>6</sup> School of Pharmacy, Kumamoto University, Japan

<sup>7</sup> All-Russian Research Institute of Phytopathology, Moscow Region, Russia

<sup>8</sup> Laboratory of Biocontrol and Antimicrobial Resistance, Orel State

University named after I.S. Turgenyev, Orel, Russia

<sup>9</sup> European Society of Clinical Microbiology and Infectious Diseases, Basel, Switzerland

<sup>10</sup> Student Research Committee, School of Medicine, Bam University of Medical Sciences Bam, Iran

<sup>11</sup> Department of Chemistry, University of Alabama in Huntsville, AL, USA

<sup>12</sup> Aromatic Plant Research Center, Lehi, UT, USA

<sup>13</sup> Food Safety Research Center (salt), Semnan University of Medical Sciences, Iran

<sup>14</sup> Department of Chemistry, Richardson College for the Environmental Science Complex, The University of Winnipeg, MB, Canada

## Corresponding Authors:

Abhay P. Mishra, Department of Pharmaceutical Chemistry, H. N. B. Garhwal University, Srinagar-246174, India.

Email: [abhaypharmachemhnbgu@gmail.com](mailto:abhaypharmachemhnbgu@gmail.com)

Bahare Salehi, Student Research Committee, School of Medicine, Bam University of Medical Sciences, Bam, Iran.

Email: [bahar.salehi007@gmail.com](mailto:bahar.salehi007@gmail.com)

William N. Setzer, Department of Chemistry, University of Alabama in Huntsville, AL, USA.

Email: [setzerw@uah.edu](mailto:setzerw@uah.edu)

Javad Sharifi-Rad, Zabol Medicinal Plants Research Center, Zabol University of Medical Sciences, Zabol, Iran.

Email: [javad.sharifirad@gmail.com](mailto:javad.sharifirad@gmail.com)



of volatile components such as terpenoids, phenylpropanoids, and aliphatic compounds.<sup>12</sup>

The 500 species of *Artemisia* are mostly perennial herbs dominating the vast steppe communities of Asia. Asia has the greatest concentration of species, with 150 accessions for China, 174 in the former Soviet Union, about 50 reported for Japan, 35 species of the genus found in Iran, and about 30 in Italy. *Artemisia* species are frequently utilized for the treatment of different diseases such as malaria, hepatitis, cancer, inflammation, and infections by fungi, bacteria, and viruses.<sup>2,5</sup>

## Traditional and Current Uses

One of the most known plants of the genus *Artemisia* is *A. absinthium* L., commonly known as “wormwood,” a yellow-flowering perennial plant distributed throughout various parts of Europe, the Middle East, North Africa, and Asia, and several chemotypes have been recognized.<sup>13</sup> The plant is used for its antiparasitic effects and to treat gastrointestinal problems, anorexia, and indigestion.<sup>14</sup> The aerial parts are present in many gastric herbal preparations, in dietary supplements, and in alcoholic beverages, for example, absinthe products, which enjoy a resurgence of popularity all over the world.<sup>2,15</sup> Moreover *A. absinthium* and other plants of this genus were used to control pain in childbirth and to induce abortions.<sup>5-17</sup>

In North African and Middle Eastern countries, *A. abyssinica* Sch. Bip. ex A. Rich. is used in folk medicine as an anthelmintic, antispasmodic, antirheumatic, and antibacterial agent.<sup>18</sup> This plant grows abundantly in various parts of the Arabian peninsula and is locally known as “ather” (Saudi Arabia) and “boitheran” (Yemen).<sup>2,19,20</sup>

Preparations of *A. abrotanum* L. (“southernwood”) have been used in traditional medicine for treating a variety of disorders, including upper airway diseases. Moreover it has been found to possess spasmolytic activity on the carbacholine-induced contraction of guinea pig trachea.<sup>6,21</sup> Nowadays, this perennial plant is used mainly for culinary or cosmetic purposes.<sup>2</sup>

*Artemisia afra* Jacq. ex Willd. is a well-known medicinal plant of South Africa, where it is known as “wilde als.” It is widely used for numerous ailments including colds, coughs, diabetes, heartburn, bronchitis, and asthma.<sup>2,22</sup>

*Artemisia annua* L. (“sweet wormwood,” “qinghao”) has traditionally been used in China for the treatment of fever and chills. Though originally growing in Asia and Europe, the plant is cultivated in Africa and used as a tea for the treatment of malaria. Artemisinin has been identified as the antimalarial principal of the plant, and artemisinin derivatives are currently established as antimalarial drugs with activity toward otherwise drug-resistant *Plasmodium* infections.<sup>2,23</sup> Most importantly, however, *A. annua* is now known worldwide for its antimalarial properties. Other *Artemisia* species have also been used for the treatment of fevers and malaria. *Artemisia absinthium* and *A. abrotanum* were used to

treat malaria in Europe, while *A. afra* in Africa.<sup>5,16,24,25</sup> The species *A. annua* and *A. apiacea* Hance are native to China. There has been some confusion about their ancient Chinese names. In older texts, *qing bao* (blue-green herb) and *cao bao* (herbaceous herb) were used interchangeably. The polymath Shen Gua (1031–1095) of the Song dynasty described two different varieties of *qing bao*, one with blue-green leaves, the other with yellowish-green leaves in autumn. Based partly on his description, the famous physician and natural historian Li Shizhen (1518–1593), whose cyclopedic *Classified Materia Medica* (*Ben cao gang mu*) was published posthumously in 1596, differentiated between *qing bao* (blue-green herb) and *huang bua bao* (yellow blossom herb).<sup>5,26</sup>

*Artemisia annua* has been recognized as an important ethnomedicinal herb for 2 millennia. It has been included in ancient pharmacopeias of various Asian and European countries. The World Health Organization has recommended *A. annua* as an antimalarial drug.<sup>27</sup> Its most common ethnobotanical practice involves the use of whole plant decoction for the treatment of malaria, cough, and cold. Its dry leaf powder has been reported in the treatment of diarrhea.<sup>28</sup> The whole flowering plant is known to be anthelmintic, antipyretic, antiseptic, antispasmodic, carminative, stimulant, tonic, and stomachic. The tincture was formally used to treat nervous diseases and crushed plants in liniments.<sup>28</sup> *Artemisia annua* tea infusion has been used for the treatment of malaria in African countries. As mentioned above, *A. annua* contains artemisinin, which provides a structural chemical base for combinatorial treatment therapy for worldwide antimalarial programs. Research studies also report that artemisinin is effective for killing human breast cancer cells.<sup>28</sup> Therefore, isolation and characterization of artemisinin has increased the interest in *A. annua* worldwide. Several ethnobotanical uses in Africa claim that the *A. annua* tea is also effective against human immunodeficiency virus (HIV). Recently, research investigations are more focused to evaluate its antiviral potential against HIV, as it is a highly emerging disease throughout the world.<sup>29</sup>

*Artemisia arborescens* L. (“great mugwort,” “arborescent mugwort”) is a morphologically variable species (or mixture of species) with grey-green to silver leaves. It is native to the various habitats of the Mediterranean region, where it occurs as a shrub growing up to 1 m in height. According to popular folklore, it is used as an anti-inflammatory remedy.<sup>2,30</sup>

*Artemisia argyi* H. Lév. & Vaniot is an herbaceous perennial plant with a creeping rhizome. It is native to China, Japan, and the far eastern parts of the former Soviet Union. In Japan, it is known as “gaiyou” and in China as “ai ye.” It is used in herbal medicine for pathologic conditions of the liver, spleen, and kidney.<sup>2</sup>

The powdered leaves of *Artemisia biennis* Willd. are used as spices and in folk remedies as antiseptics. They have been applied externally in salves and washes by the native

**Table 1.** List of the Chemical Constituents of Various *Artemisia* Spp.

| S. no. | <i>Artemisia</i> species         | Chemical constituents   | References |
|--------|----------------------------------|---|------------|
| 1.     | <i>A. annua</i>                  | Artemisinin, scopoletin, arteannuin B and arteannuic acid, 5- <i>O</i> -[( <i>E</i> )-caffeoyl]quinic acid, 1,3-di- <i>O</i> -caffeoylquinic acid, 4,5-di- <i>O</i> -caffeoylquinic acid, 3,5-di- <i>O</i> -caffeoylquinic acid, 3,4-di- <i>O</i> -caffeoylquinic acid, methyl-3,4-di- <i>O</i> -caffeoylquinic acid, methyl-3,5-di- <i>O</i> -caffeoylquinic acid, 3,6'- <i>O</i> -diferuloylsucrose, 5'- $\beta$ -D-glucopyranosyloxyljasmonic acid, scoparone, 4- <i>O</i> - $\beta$ -D-glucopyranosyl-2-hydroxyl-6-methoxyacetophenone, chrysosplenol D, casticin, chrysosplenetin, artemisinic acid, deoxy-artemisinin, artemetin, 7,8-dimethylalloxazine (lumichrome), daucosterol, <i>p</i> -hydroxybenzoic acid, uracil, nicotinic acid, 3 $\alpha$ -hydroxy-1-deoxyartemisinin, salicylic acid, domesticoside (2- <i>O</i> - $\beta$ -D-glucopyranosyl-4- <i>O</i> -methylphloroacetophenone, scopolin, $\beta$ -sitosterol, quercetagenin-6,7,3',4'-tetramethyl ether, quercetagenin-6,7,4'-trimethyl ether                             | 73-77      |
| 2.     | <i>A. judaica</i>                | Oxygenated monoterpenes, sesquiterpenes, and hydrocarbons   | 78         |
| 3.     | <i>A. herba-alba</i>             | Oxygenated monoterpenes, sesquiterpenes and hydrocarbons, chlorogenic acid, 4,5- <i>O</i> -dicaffeoylquinic acid, isofraxidin 7- <i>O</i> - $\beta$ -D-glucopyranoside, 4- <i>O</i> - $\beta$ -D-glucopyranosylcaffeic acid, rutin, schaftoside, isoschaftoside, and vicenin-2  | 78         |
| 4.     | <i>A. vulgaris</i>               | Eudesmane-type sesquiterpene, morin, luteolin, triterpenes, coumarin, flavonoids, eriodictyol   | 79-81      |
| 5.     | <i>A. rupestris</i>              | Citrusin A, alachanioside A, coniferin, citrusin B, syringaresinol- $\beta$ -D-glucoside, (6 <i>R</i> ,9 <i>S</i> )-3-carbonyl- $\alpha$ -ionolglucopyranoside, byzantioside B, 6-demethoxy-4'- <i>O</i> -methylcapillarisin-7- <i>O</i> - $\beta$ -D-glucopyranoside, catechin, kaempferide, artemetin, kaempferol 3,3',4'-trimethylether, umbelliferone, stigmaterol, rupestonic acid, artemetin, casticin, chlorogenic acid, 7-hydroxycoumarin, rupestonic acid, 12-hydroxy-4,11(13)-guaiadien-3-one, luteolin, 5,7-dihydroxy-3',4'-dimethoxyflavone, luteolin 7- <i>O</i> - $\beta$ -D-glucoside, luteolin 7- <i>O</i> -[ $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside], linarin, quercetin, nevadensin, gardenin D, skimmin, stigmaterol-3- <i>O</i> - $\beta$ -D-glucopyranoside, 1,2-isopropylidene- $\alpha$ -D-glucopyranose, tianshic acid, $\beta$ -sitosterol, daucosterol, rupestonic acid, chrysosplenetin B, herniarin, isokaempferide, vanillic acid, kaempferol 3,3',4'-trimethyl ether, and ermanin | 82-84      |
| 6.     | <i>A. capillaris</i>             | Neochlorogenic acid, chlorogenic acid, cryptochlorogenic acid, caffeic acid, 1,3-dicaffeoylquinic acid, 3,4-dicaffeoylquinic acid, 3,5-dicaffeoylquinic acid, 4,5-dicaffeoylquinic acid, chlorogenic acid analogs and phenolic acids, 7-hydroxy-coumarin, 5,7-dimethoxy-coumarin, 7-hydroxy-8-methoxy-coumarin, 7,8-dihydroxy-coumarin, quercetin, kaempferol, isorhamnetin-3- <i>O</i> - $\beta$ -D-galactopyranoside, 7-methoxycoumarin, 4-hydroxyacetophenone, nicotinic acid, 3( <i>R</i> )-deca-4,6,8-triyn-1,3-diol, 3( <i>R</i> )-deca-4,6,8-triyn-1,3-diol-1- <i>O</i> - $\beta$ -D-glucopyranoside, 3( <i>R</i> )-9-decene-4,6-diyne-1,3,8-triol, 4-hydroxyacetophenone-4- <i>O</i> - $\beta$ -D-glucopyranoside, phenylcarbinol- <i>O</i> - $\beta$ -D-glucopyranoside, thymidine, isoquercitrin, isorhamnetin-3- <i>O</i> -glucoside, ursolic acid, oleanolic acid, and $\beta$ -sitosterol  | 85-89      |
| 7.     | <i>A. ordosica</i> Krasch        | 3,5,3',4'-Tetrahydroxy-6,7-dimethoxyflavone, 5,3',4'-trihydroxy-7-methoxyflavanone, 5,7,4'-trihydroxy-6-methoxyflavone, 5,7,4'-trihydroxyflavanone, 3,5,4'-trihydroxy-7-methoxyflavanone, 3,5,7-trihydroxy-4'-methoxyflavanone, isosakuranetin, 7,4'-dimethylaromadendrin, acetatin, cirsimaritin, rhamnetin, eupatolitin, 5,7,2',4'-tetrahydroxy-6,5'-dimethoxyflavone, hyperoside   | 90,91      |
| 8.     | <i>A. absinthium</i>             | Absinthin, anabsinthin, 5,6,3',5'-tetramethoxy-7,4'-hydroxyflavone, 5-hydroxy-3,3',4',6,7-pentamethoxyflavone, artemetin, rutin, glycosides of quercetin, chlorogenic, caffeic acids, artamarin, artamaridin, artamaridin, artamarinin, quebrachitol  | 92-94      |
| 9.     | <i>A. sphaerocephala</i> Krasch. | Caffeic acid, 5,7,2',4'-tetrahydroxy-6,5'-dimethoxyflavone, quercetagenin, 4'-hydroxywogonin, quercetagenin-4'-methyl ether, 5,6,3'-trihydroxy-7,4'-dimethoxyflavanone, eriodictyol-7-methyl ether, quercetin-7-methyl ether, sakuranetin, isosakuranetin, 3,5-dihydroxy-7,4'-dimethoxy-flavanone, 5-hydroxy-7,4'-dimethoxyflavanonol, 5,3'-dihydroxy-7,4'-dimethoxyflavanone, 5,7-dihydroxy-6,4'-dimethoxyflavone, hesperetin, naringenin, acetatin, chrysoeriol, 5,7-dihydroxy-4'-methoxyflavone-6,8-di- <i>C</i> -glucopyranoside, didymin, acetatin-7- <i>O</i> -rutoside, piceine, capillarin, ethyl linoleate, sakuranetin, isosakuranetin, artocarpanone   | 90,95,96   |
| 10.    | <i>A. turanica</i>               | 3,5-Dicaffeoylquinic acid, 4,5-dicaffeoylquinic acid, 3,5,3',4'-tetrahydroxy and 7,5'-methoxy flavones  | 97,98      |
| 11.    | <i>A. campestris</i>             | Catechin, vanillic acid, caffeic acid, syringic acid, <i>p</i> -coumaric acid, and gallic acid  | 99         |
| 12.    | <i>A. gmelinii</i>               | Genkwanin, hispidulin, 3'-hydroxy-genkwanin, chrysoeriol, apigenin, 5,7,3',4'-tetrahydroxy-6,5'-dimethoxy-flavone, kumatakenin, quercetin, patuletin, quercetagenin-3,6,7-trimethylether, and 7,3',4'-trihydroxy-3-methoxy-flavone  | 100        |

(Continued)

Table 1. Continued

| S. no. | <i>Artemisia</i> species | Chemical constituents  | References |
|--------|--------------------------|--|------------|
| 13.    | <i>A. selengensis</i>    | ( <i>E</i> )-Resveratrol, ( <i>E</i> )-cinnamic acid, caffeic acid, chlorogenic acid, gallic acid, luteolin, isorhamnetin, 7-methoxy coumarin, quercetin, acteoside, 7-methoxy-4'-hydroxyisoflavone, 1',3'-propanediol,2'-amino-1'-(1,3-benzodioxol-5-yl), artanomaloid, canin, eupatilin, quercetin-3- <i>O</i> - $\beta$ -D-glucoside-7- <i>O</i> - $\alpha$ -L-rhamnoside, isoquercitrin, 1,3-di- <i>O</i> -caffeoylquinic acid, pinoresinol-4- <i>O</i> - $\beta$ -D-glucoside, scopolin, isofraxidin-7- <i>O</i> - $\beta$ -D-glucopyranoside, $\beta$ -farnesene, <i>N</i> -( $\alpha$ -methylnaphthalene)-3- ( <i>ortho</i> -cresyl) propionamide, <i>N</i> -( $\alpha$ -methylnaphthalene)-3-( <i>contra</i> -cresyl) propionamide, $\alpha$ -pinene oxide, $\alpha$ -humulene, $\alpha$ -cedrene, $\beta$ -curcumene, zingiberene, sesquisabinene hydrate, isocaryophyllene, $\beta$ -sesquiphellandrene, 4',7-dimethyluteolin, 11,13-dihydromatricarin, chrysoerid-7- <i>O</i> - $\beta$ -D-glucoside, luteolin-4'- <i>O</i> - $\beta$ -D-glucoside, quercetin-3- <i>O</i> - $\beta$ -D-xyloside, daucosterol  | 101–105    |
| 14.    | <i>A. scoparia</i>       | 4-Pyridone glucoside, polyacetylene glucosides, 2-(5-acetyl-2,3-dihydro-benzofuran-2-yl)-propyl ester, 2-(3'-methoxyl-3'-methyl- <i>trans</i> -but-1'-enyl)- <i>p</i> -methoxyacetophenone, 5,8-dimethoxy-6,7-methylenedioxy coumarin (Artemicapin A), 8-methoxy-6,7-methylenedioxy coumarin, 6-demethoxycapillarisin, 2,4-dihydroxyl-6-methoxyacetophenone, $\beta$ -sitosterol, 6,7- dimethoxy-coumarin, capillartemisin B, kumatakenin and isoscopoletin- $\beta$ -D-glucoside, 7-methoxycoumarine, isosabandin, 6,7-dimethylesculetin, 7-methylesculetin, scopoletin, capillarisin, chlorogenic acid butyl ester, isoscopoletin- $\beta$ -D-glucoside (magnolioside), hyperin, eupafolin, pedaltin, 5,7,2',4'-tetrahydroxy-6,5'-dimethoxyflavone, capillone, capillin, jaceosidin, chrysoeriol, arcapillin, luteolin, chrysoeriol-7- <i>O</i> - $\beta$ -D-glucopyranoside, cacticin, isorhamnetin-3- <i>O</i> - $\beta$ -D-glucopyranoside, quercetin-7- <i>O</i> - $\alpha$ -L-rhamnopyranoside, hyperin and <i>n</i> -butyl- $\beta$ -D-fructopyranoside, cirsilineol, cirsimaritin, arcapillin, cirsilio, quercetin-3- <i>O</i> - $\beta$ -D-glucoside | 86,106-111 |
| 15.    | <i>A. frigida</i>        | 3,5-dihydroxy-5-methoxycinnamic acid, caffeic acid, 6,7-dimethoxycoumarin  | 112        |
| 16.    | <i>A. lactiflora</i>     | Kaempferol, ferulic acid, cinnamic acid, daucosterol, 5,7,3',4'-tetrahydroxy-6,5'-dimethoxyflavone, 5,7-dihydroxy-6,3',4'-trimethoxyflavone, 5,7,4'-trihydroxy-6,3'-dimethoxyflavone, 5,7,4'-trihydroxy-6-methoxyflavone, 5,7,3',4'-tetrahydroxy-6-methoxyflavone, rutin , 3'-methoxy-luteolin-4'- <i>O</i> - $\beta$ -D-glucoside, 5-hydroxy-3',4'-dimethoxyflavone-7- <i>O</i> - $\beta$ -D-glucuronide  | 113-115    |
| 17.    | <i>A. indica</i>         | 7-Hydroxycoumarin, 7-methoxycoumarin, balanophonin, aurantiamide, aurantiamide acetate, isovitexin, kaempferol-3- <i>O</i> - $\beta$ -D-rutinoside, rutin, caffeic acid ethyl ester, quercetin, methyl 3,5-di- <i>O</i> -caffeoyl quinate, methyl 3,4-di- <i>O</i> -caffeoyl quinate, dehydromifoliol, 3-hydroxy-1-(4-hydroxy-3,5-dimethoxy-phenyl)-1-propanone, chrysindin D, camelliagenin A, 4'- <i>O</i> -methylalpinumisoflavone, 5-hydroxy-3',4',6,7,8-pentamethoxyflavone, armexifolin, 3 $\beta$ -hydroxy-5 $\alpha$ ,6 $\alpha$ -epoxy-7-megastigmen-9-one, carissone, and ( <i>E</i> )-3 $\beta$ ,4 $\alpha$ - dihydroxyl-(2',4'-hexadiynylidene)-1,6-dioxaspiro [4,5] decane, 4-(2-hydroxyethoxy) acetophenone, loliolide, isolololide, isovanillic acid, <i>p</i> -hydroxybenzoic acid, ( <i>E</i> )- <i>p</i> -coumaric acid, ethyl ( <i>E</i> )- <i>p</i> -coumarate, (+)-pinoresinol, (+)-medioresinol, (+)-syringaresinol, 4-hydroxy-3,5-dimethoxy-benzoic acid, dihydroisoferulic acid  | 116        |
| 18.    | <i>A. leucophylla</i>    | 5-Hydroxy-3,7,4'-trimethoxyflavone, ludartin, maackiain, lupeol, <i>cis</i> -matricaria ester, <i>trans</i> -matricaria ester, and 6-methoxy-7,8-methylenedioxy coumarin   | 117        |
| 19.    | <i>A. myriantha</i>      | Lupeol, apigenin, luteolin, isorhamnetin, monoheptadecanoin, scopoletin, umbelliferone, kaempferol-3- <i>O</i> - $\alpha$ -LL-rhamnoside, kaempferol-3- <i>O</i> -glucoside, kaempferitrin and D-mannitol  | 74         |
|        |                          | Arglabin, 13-acetoxy-3 $\beta$ -hydroxyl-germacra- 1(10) <i>E</i> ,4 <i>E</i> ,7(11)-trien-12,6 $\alpha$ -olide, eupatorin, 8 $\alpha$ -acetoxyarglabin, artemyriantholide B, artemyriantholide A, 4',5,7-trihydroxy-6,3'-dimethoxy flavone, cacticin, 5,4'-dihydroxy-6,7,3',5'-tetramethoxyflavone, arborescin, arlatin   |            |

(Continued)

Table 1. Continued

| S. no. | <i>Artemisia</i> species  | Chemical constituents  | References     |
|--------|---------------------------|--|----------------|
| 20.    | <i>A. anomala</i>         | $\beta$ -Sitosterol, $\beta$ -daucosterol, schleicheol 2, $\alpha$ -spinasterol, 5 $\alpha$ ,8 $\alpha$ -epidioxy-ergosta-6,22-dien-3 $\beta$ -ol, 5 $\alpha$ ,8 $\alpha$ -epidioxy-ergosta-6,9(11),22-trien-3 $\beta$ -ol, 22E-3 $\beta$ ,5 $\alpha$ -dihydroxyergosta-7,22-dien-6-one, naringenin, luteolin, kaempferol, chrysoeriol, diosmetin, jaceosidin, isorhamnetin 3-O-glucoside, hesperetin-7-O- $\beta$ -D-glucopyranoside, methyl (4a <i>S</i> ,7 <i>S</i> ,7a <i>R</i> )-7-hydroxy-7-methyl-1,4a,5,6,7,7a-hexahydrocyclopenta[c]pyran-4-carboxylate, rehmaglutin D, (E)-6-hydroxy-2,6-dimethylocta-2,7-dienoic acid, chrysoeriol, luteolin, apigenin, <i>p</i> -coumaric acid, 3 $\beta$ -ethoxytanaparatholide, (4 <i>S</i> *,5 <i>S</i> *)-dihydro-5-[(1 <i>R</i> *,2 <i>S</i> *)-2-hydroxy-2-methyl-5-oxo-3-cyclopenten-1-yl]-3-methylene-4-(3-oxobutyl)-2(3H)-furanone, ligucyperonol, cyperusol C, santamarin, 1 $\alpha$ ,2 $\alpha$ ,3 $\alpha$ ,4 $\alpha$ ,10 $\alpha$ -pentahydroxyguaia-11(13)-ene-12,6 $\alpha$ -olide, balanophonin, methyl 3-(2'-hydroxy-4'-methoxyphenyl) propanoate, acceroic acid, simiarenol, $\alpha$ -amyrin, $\beta$ -sitosterol, 4-methoxysalicylic acid, <i>m</i> -hydroxybenzoic acid, 2'-hydroxycinnamaldehyde, cinnamic acid, isoferulic acid, 7-methoxycoumarin, indolyl-3-carboxylic acid, dihydrophaseic acid 4'-O- $\beta$ -D-glucopyranoside, citroside A, (6 <i>S</i> ,9 <i>R</i> )-roseoside, cyclobalanone, friedelin, sorghumol, pseudoneolinderane, herniarin, scopoletin, isofraxidin and caffeic acid | 118-122        |
| 21.    | <i>A. argyi</i>           | Lemaphenol A, aurantiamide acetate, camelliagenin A, japonica acid, labd-13(E)-ene-8 $\alpha$ ,15-diol, 3 $\beta$ -acetoxy-20-oxo-21-nordammaran-23-oic acid, apigenin, jaceosidin, luteolin, eupatilin, $\beta$ -sitosterol, quercetin, isotanciloide, umbelliferone, daphnetin, eriodictyol, rhamnetin, hispidulin, 1-2-O-methyl-chiro-inositol, stigmasterol, daucosterol, and 4-methoxy-3-hydroxyphenol  | 123,124<br>125 |
| 22.    | <i>A. rupestris</i>       | 5,4'-Dihydroxy-3,6,7-trimethoxy flavone, R(-)-vestitol, tricin, chrysoeriol, 3-indole carboxylic acid, esculetin, apigenin, luteolin, luteolin-7-glucoside, (E)-caffeic acid, casticin, chrysopterin B, artemetin, robinin, quercetin, linearin, sucrose, tilianin, 3'-methoxy-4'-O- <i>p</i> -coumaroyl- $\beta$ -D-glucoside, hirsutine, ethyl <i>p</i> -methoxycinnamate, rutin   | 126,127        |
| 23.    | <i>A. sacrorum</i>        | $\beta$ -Sitosterol, acacetin, 7-methoxy-6-hydroxycoumarin, 5-hydroxy-7,4'-dimethoxyflavone, 1,4-dicaffeoylquinic acid, salicylic acid, veratric acid, scopoletin, isofraxidin, succinic acid, sugereoside, sacroside C, <i>o</i> -hydroxycinnamoyl- $\beta$ -D-glucopyranoside, 6-methoxycoumarin-7-hydroxylprimeveraside   | 128,129        |
| 24.    | <i>A. amygdalina</i>      | Ergostadien-3 $\beta$ -ol, ludartin, 5-hydroxy-6,7,3',4'-tetramethoxyflavone, <i>trans</i> -matricaria ester, diacetylenic spiroenol ether, and <i>cis</i> -matricaria ester   | 130            |
| 25.    | <i>A. lavandulaefolia</i> | Kaempferol, formononetin, isorhamnetin, apigenin, tricin, quercetin-7-O- $\beta$ -D-glucopyranoside, quercetin-7-O- $\alpha$ -L-rhamnopyranoside, quercetin-3-O- $\beta$ -D-glucopyranoside, kaempferol-3-O- $\beta$ -D-glucopyranoside, apigenin-7-O- $\beta$ -D-glucopyranoside, kaempferol-3-O- $\beta$ -D-rutinoside, 7-methoxycoumarin, <i>m</i> -hydroxybenzoic acid, isoferulic acid, lolilide, medioresinol, (+)-syringaresinol, syringic acid, friedelinol, caffeic acid, vanillic acid, stigmasterol, sitosterol-3-O-glucopyranoside, $\beta$ -amyrin, $\beta$ -sitosterol, ursolic acid, eupatilin, naringenin, apigenin, luteolin, quercetin   | 131,132        |
| 26.    | <i>A. dubia</i>           | Calotropoleanyl ester, $\alpha$ -amyrin, nonacosanoic acid, docosanoic acid, tetracosanoic acid, 1-(O-tricosanoyl) glycerol, 1-(O-pentacosanoyl) glycerol, and $\beta$ -sitosterol   | 133            |
| 27.    | <i>A. dracunculus</i>     | 1-Monopalmitin, methyl $\alpha$ -linolenate, 1-monoolein, 3-O-acetylursolic acid, 3 $\beta$ ,22 $\beta$ ,24-trihydroxy-olean-12-ene, oleic acid, palmitic acid, 1-monolinolein, vanillic acid, hesperetin, naringenin, quercetin, rutin, scoparone, scopoletin, 7-hydroxycoumarin, daphnetin, caffeic acid, chlorogenic acid, 7-methoxycoumarin, 7-hydroxyartemidin  | 134,135        |
| 28.    | <i>A. inayomogi</i>       | Coumarins, phenolics, flavonoids, caffeoylquinic acids, diterpene glycosides, hispidulin, 6-methoxytricin, arteanoflavone, quercetin-3-gentiobioside, 1,3-di-O-caffeoylquinic acid, suavioside A, turpinionoside A, (Z)-3-hexenyl-O- $\alpha$ -arabinopyranosyl-(1 $\rightarrow$ 6)-O- $\beta$ -D-glucopyranoside, (Z)-5'-hydroxyjasmonone 5'-O- $\beta$ -D-glucopyranoside, (-)-syringaresinol-4-O- $\beta$ -D-glucopyranoside, and methyl 3,5-di-O-caffeoyl quinate  | 136,137        |
| 29.    | <i>A. balodendron</i>     | Arcapillin, isorhamnetin-3-O-glucopyranoside, <i>p</i> -hydroxyacetophenone, caffeic acid, <i>p</i> -coumaric acid, isorhamnetin, palmitic acid, cirsilineol, pentacosanol   | 138            |
| 30.    | <i>A. waldst</i>          | 1-Dotriacontanol, palmitic acid, hexacosyl eicosanoate, $\beta$ -daucosterol   | 139            |
| 31.    | <i>A. inculta</i>         | Germacrene D, phytol, santolin alcohol, a hydroperoxide, 4,5-dihydroxysantolin-1,8-diene, a sesquiterpene lactone, flavone artemetin   | 140            |
| 32.    | <i>A. vestita</i>         | Taurin, 1,11-bis- <i>epi</i> -artemin, yomogin, 5,7,3',4'-tetrahydroxy-6, 8-dimethoxy flavone, 5,7,3',4'-tetrahydroxy-6-methoxy flavone, 5,6,3',4'-tetrahydroxy-7-methoxy flavone, taraxerol-3 $\beta$ -acetate, friedelin, $\alpha$ -amyrin, $\beta$ -sitosterol, daucosterol, scoplatin, 7-hydroxy-6,8-dimethoxycoumarin, isoferulic acid, caffeic acid  | 141            |

(Continued)

Table 1. Continued

| S. no. | <i>Artemisia</i> species | Chemical constituents  | References |
|--------|--------------------------|--|------------|
| 33.    | <i>A. arborescens</i>    | Artemitin, arborescin, sesamin, (+)-lirioresinol $\beta$ -dimethyl ether, chrysoeriol, apigenin, $\beta$ -sitosteryl glucoside, dihydridentin, and chrysoeriol 4-glucoside   | 142        |
| 34.    | <i>A. roxburgiana</i>    | Riedelin, $\alpha$ -amyrin acetate, $\alpha$ -amyrin, $\beta$ -amyrin acetate, $\beta$ -sitosterol, dotriacontanoic acid, octacosanol, octacosanoic acid, triacontanoic acid, daucosterol, friedelan-3-one, multiflorenone, stigmastanone, 8-hydroxydotriacontan-27-one  | 143        |
| 35.    | <i>A. tangutica</i>      | $\beta$ -Sitosterol, $\alpha$ -amyrin acetate, $\alpha$ -amyrin, $\beta$ -amyrin, $\beta$ -amyrin acetate, 2-oxo-desoxyglustrin, estafiatin, 4 $\alpha$ -hydroxyguaia-10(14),11(13)-dien-12,6 $\alpha$ -olide, 1- <i>epi</i> -reynosin, and 9 $\beta$ -acetoxy-4,5-dehydro-4(15)-dihydrocostic acid  | 144        |
| 36.    | <i>A. japonica</i>       | $\beta$ -Amyrin, triacontanoic acid, $\beta$ -sitosterol, stigmasterol, 7,8-dimethoxycoumarin, 6,7-dimethoxycoumarin, capillarisin, 8,4'-dihydroxy-3,7,2'-trimethoxyflavone, 3,5-dihydroxy-6,7,3',4'-tetramethoxyflavone, cinnamic acid, <i>p</i> -methoxybenzoic acid, and ferulic acid   | 145        |
| 37.    | <i>A. subdigitata</i>    | 5,8,3',5'-Tetrahydroxyflavanone, 5,8,2'-trihydroxy-5'-methoxyflavanone, 5,7,4'-trihydroxy-3',5'-dimethoxyflavanone, 3-(3-hydroxy)-phenoxy-2-propenal, tricrin, quercetin-3-rhamnoside, ethyl 2,5-dihydroxycinnamate, 8-hydroxy-6,7-dimethoxycoumarin, 2,4-hexadiyn-1-one, $\alpha$ -amyrin, $\beta$ -amyrin, $\alpha$ -amyrin acetate, $\alpha$ -amyrone, friedelin, $\beta$ -sitosterol, tetracosanoic acid, and <i>n</i> -butyl palmitate  | 146        |
| 38.    | <i>A. montana</i>        | Ezoartemin, ezomontanin, 11,13-dihydroezomontanin, yomogiartermine, yamayomoginin  | 147        |
| 39.    | <i>A. myriantha</i>      | Blumenol A, (+)-dehydrovomifoliol, (+)-3-hydroxy- $\beta$ -ionone, (3R,6R,7E)-3-hydroxy-4,7-megastigmadien-9-one, (-)-10-oxo-isodauc-3-en-15-oic acid, isoerivanin, eudesmafraglaucolide, artanomalide A, 13-acetoxy-3 $\beta$ -hydroxy-germacra-1E,4E,7(11)-trien-12,6 $\alpha$ -olide, 13-acetoxy-3 $\beta$ -tigloyl-germacra-1E,4E,7(11)-trien-12,6 $\alpha$ -olide, 13-acetoxy-3 $\beta$ -(3-methylbutanoyl)-germacra-1E,4E,7(11)-trien-12,6 $\alpha$ -olide, 3,9-diacetoxy-13-hydroxy-1(10),4,7(11)-germacatrien-12,6 $\alpha$ -olide, and 8 $\alpha$ -angeloyloxycostunolide | 148        |

inhabitants of North America for treating sores and wounds, and internally to treat chest infections.<sup>2,31</sup>

*Artemisia campestris* L. is a perennial faintly aromatic herb, widespread in the south of Tunisia, commonly known as “tgouft.” The leaves of this plant are widely used in traditional medicine as a decoction for their antivenin, anti-inflammatory, anti-rheumatic, and antimicrobial properties.<sup>32</sup> *Artemisia cana* Pursh is used as a spice and in folk remedies as an antiseptic.<sup>2</sup>

*Artemisia douglasiana* Besser (“California mugwort”) is a perennial herb that is native to the western United States, especially northern California, Oregon, and Washington. *Artemisia douglasiana* is used to promote menstruation, as a stimulant, tonic, to treat nervous disorders, and as a diuretic. The essential oil has been used for aromatherapy, inhaled for mental clarity and ease of mental distress; used as a massage for aching muscles and pain on the surface of the body; and as a bath or tonic.<sup>33</sup> In Argentina, *A. douglasiana*, which is adventitious and cultivated in the Cuyo region, is used in folk medicine and known under the common name of “matico.” The popular use of the infusion of leaves of “matico” is to treat peptic ulcers and gastrointestinal disorders.<sup>34</sup>

*Artemisia dracunculoides* L. (“tarragon”) is a perennial herb, which has a long history of use in culinary traditions. It also possesses a wide range of health benefits and has therefore been widely used as an herbal medicine. For example, in the Himalayas, extracts of *A. dracunculoides* are used to relieve toothache, reduce fever, and as a treatment for gastrointestinal problems.<sup>35</sup> Two well-described cultivars (Russian and French) are used widely and differ in ploidy level, morphology, and chemistry. The botanical and chemical constituents are

closely detailed in the literature, the latter mainly focusing on its essential oil composition, which give its distinctive flavor.<sup>2,36</sup>

*Artemisia dubia* Wall. ex Besser is native to Bhutan, China, India, Japan, Nepal, and Thailand.<sup>37</sup> In Nepal, the leaf juice of *A. dubia* is used to treat cuts and wounds while the plant paste is used against fever.<sup>38</sup>

*Artemisia ecbegaray* Hieron is commonly known in Argentina as “ajeno” and is used as a natural food additive.<sup>2</sup> Decoctions of leaves and stems of *A. frigida* Willd. are used for coughs and diabetes<sup>2</sup>; several native American tribes have used decoctions of *A. frigida* for menstrual irregularities.<sup>39</sup>

*Artemisia fukudo* Makino is distributed along the shorelines of South Korea's Jeju Island and in the south of the Korean Peninsula, Japan, and Taiwan. This plant is used as a flavoring agent and in a variety of cosmetics in Korea. It also has various biological effects, including anti-inflammatory, antitumor, and antibacterial properties.<sup>40</sup>

*Artemisia gmelinii* Weber ex Stechm. is a plant of the high-altitude regions of Asia. In Nepal, the fresh plant is ground into a paste and applied externally for headaches, boils, and pimples.<sup>35</sup> *Artemisia haussknechtii* Boiss. is used in dyspepsia and other gastrointestinal disorders by local people in the Western part of Iran (province of Kermanshah).<sup>41</sup> In Nepal, *A. indica* Willd. is used to treat ringworm, cuts, and wounds, and as an anti-leech treatment.<sup>35,38</sup> *Artemisia inayomogi* Kitamura is a perennial herb easily found around Korea. It is called “hanin-jin” or “dowijigi” in Korean and is traditionally used for the treatment of various liver diseases, including hepatitis.<sup>2</sup>

**Table 2.** List of the Constituents of Essential Oils in Various *Artemisia* Spp.

| S. no. | <i>Artemisia</i> species | Essential oil components  | References        |
|--------|--------------------------|---|-------------------|
| 1.     | <i>A. giraldi</i>        | 1,8-Cineole, camphor, terpinen-4-ol, $\alpha$ -terpineol  | 149               |
| 2.     | <i>A. rubripes</i>       | $\beta$ -Farnesene, 1,8-cineole, $\beta$ -caryophyllene, germacrene D, camphor  | 149,150           |
| 3.     | <i>A. judaica</i>        | $\beta$ -Eudesmol, palmitic acid, spathulenol, eudesma-4 (15),7-dien-1- $\beta$ -ol, carvacrol, thymol, piperitone, camphor and ethyl ( <i>E</i> )-cinnamate  | 78,151            |
| 4.     | <i>A. herba-alba</i>     | Piperitone, ethyl ( <i>E</i> )-cinnamate, ethyl ( <i>Z</i> )-cinnamate, thymol, isophorone, 1,8-cineole, <i>cis</i> -pinocarveol, artemisia ketone, $\alpha$ -thujone, germacrene D, camphor and $\beta$ -thujone   | 78,152,153        |
| 5.     | <i>A. ciniformis</i>     | Camphor, 1,8-cineole and <i>trans</i> -pinocarveol  | 154               |
| 6.     | <i>A. vulgaris</i>       | Sabinene, 1,8-cineole, artemisia ketone, both thujone isomers ( $\alpha$ - and $\beta$ -diastereoisomers), camphor, <i>cis</i> -chrysanthenyl acetate, davanone, davanone B, chrysanthenone, borneol, germacrene D, $\alpha$ -pinene, menthol, $\beta$ -eudesmol, spathulenol, ( <i>Z,Z</i> )-3,5-octadiene, 2,5-octadiene, 3,4,5-trimethyl-1-hexene, pulegone, 3-methyl-2-cyclohexene-1-one, decahydro-1,1,7-trimethyl-4-methylene-1 <i>H</i> -cyclopropazulene  | 79,155,156<br>157 |
| 7.     | <i>A. asiatica</i>       | Piperitone, davanone, <i>p</i> -cymene and 1,8-cineole  | 158               |
| 8.     | <i>A. anethoides</i>     | 1,8-Cineole, 2-isopropyl-5-methyl-3-cyclohexen-1-one, terpinen-4-ol, <i>o</i> -cymene and pinocarveol   | 159               |
| 9.     | <i>A. dubia</i>          | Terpinolene, limonene, 2,5-etheno [4.2.2] propella-3,7,9-triene, isoelemicin and <i>p</i> -cymene-8-ol  | 160               |
| 10.    | <i>A. aucheri</i>        | Camphor, 1,8-cineole, verbenone, camphene, $\beta$ -myrcene, $\alpha$ -pinene   | 161,162           |
| 11.    | <i>A. nilagirica</i>     | $\alpha$ -Thujone, germacrene D, $\beta$ -thujone, $\beta$ -caryophyllene, caryophyllene oxide, borneol   | 163               |
| 12.    | <i>A. campestris</i>     | Germacrene D, $\beta$ -pinene, $\alpha$ -pinene, $\alpha$ -cadinol, limonene, $\beta$ -myrcene, falcariol, $\alpha$ -terpenyl acetate, camphor, spathulenol, camphene, limonene, and borneol  | 62,99             |
| 13.    | <i>A. absinthium</i>     | $\beta$ -Thujone, 1,8-cineole, <i>cis</i> -chrysanthenol, sabinene, camphor, isoascaridol, $\beta$ -pinene <i>o</i> -cymene, $\alpha$ -phellandrene, myrcene, <i>cis</i> -chrysanthenyl acetate, dihydrochamazulene isomer, germacrene D, linalool acetate, linalool, 2-methyl-5-(1-methyl-ethenyl)-2-cyclohexen-1-one, $\beta$ -caryophyllene, 1,2-dihydro-1,4,6-trimethyl-naphthalene, elemol, 6-methyl-2,2-dipyridine <i>N</i> -oxide  | 13,164-168        |
| 14.    | <i>A. dracunculus</i>    | 1,8-Cineole, camphor, camphene, borneol, thymene, terpinen-4-ol, $\gamma$ -terpinene, $\alpha$ -terpineol, caryophyllene oxide, $\beta$ -pinene, $\alpha$ -pinene, $\beta$ -myrcene, limonene, ( <i>Z</i> )- $\beta$ -ocimene, ( <i>E</i> )- $\beta$ -ocimene, $\alpha$ -terpinolene, 5-phenyl-1,3-pentadiyne, methyleugenol, capillene, elemicin, bicyclogermacrene, iso-elemicin, germacrene B, <i>p</i> -cymene, linalool, $\beta$ -thujone, camphor, methylchavicol, iso-menthol, bornyl acetate, carvacrol, $\alpha$ -terpinyl acetate, hexyl hexanoate, $\alpha$ -copaene, $\beta$ -caryophyllene, $\beta$ -phellandrene, 3,7-dimethyl-1,3,7-octatriene, 1 <i>S</i> - $\alpha$ -pinene, 1-methoxy-4-(2-propenyl)-benzene, limonene, 1 <i>R</i> - $\alpha$ -pinene | 169-171           |
| 15.    | <i>A. annua</i>          | Selin-3,11-dien-6 $\alpha$ -ol, artemisia ketone, $\alpha$ -pinene, 1,8-cineole, camphor, bisabolol, bisabolol oxide B, bisabolol oxide A, ( <i>E</i> )-nerolidol, L-borneol, $\alpha$ -copaene, $\beta$ -caryophyllene, $\beta$ -bisabolene, germacrene D, germacrene B, (-)-neoclovene-(II), isoaromadendrene epoxide, <i>cis</i> -lanceol, caryophyllene oxide, ( <i>E</i> )- $\beta$ -farnesene, 2,5-dihydro-3-methylfuran, $\beta$ -myrcene, ( <i>Z</i> )-caryophyllene, santolina triene  | 166,172-176       |
| 16.    | <i>A. montana</i>        | <i>n</i> -Palmitic acid   | 166               |
| 17.    | <i>A. millefolium</i>    | Oxygenated monoterpenes   | 166               |
| 18.    | <i>A. arborescens</i>    | $\alpha$ -Thujone   | 177               |
| 19.    | <i>A. mongolica</i>      | 1,8-Cineole, ( <i>S</i> )- <i>cis</i> -verbenol, terpinen-4-ol, (-)-camphor, $\alpha$ -terpineol, and verbenol  | 178               |
| 20.    | <i>A. capillaris</i>     | $\alpha$ -Pinene, $\beta$ -pinene, limonene, 1,8-cineole, piperitone, $\beta$ -caryophyllene, capillin, palmitic acid, 9,12,15-octadecatrienoic acid, falcariol, <i>trans</i> -( <i>Z</i> )- $\alpha$ -bisabolene epoxide, and germacrene D   | 179,180           |
| 21.    | <i>A. diffusa</i>        | Camphor, 1,8-cineole, and $\beta$ -thujone  | 181               |
| 22.    | <i>A. rupestris</i>      | $\alpha$ -Terpinyl acetate, spathulenol, $\alpha$ -terpineol, linalool, terpinen-4-ol, $\beta$ -elemene, $\beta$ -sesquiphellandrene, $\alpha$ -guaiene, linalyl 3-methyl-butyrate, 1-hexadecanol, palmitic acid, linoleic acid   | 182,183           |
| 23.    | <i>A. scoparia</i>       | Palmitic acid, caryophyllene oxide, spathulenol, $\beta$ -myrcene, (+)-limonene, ( <i>Z</i> )- $\beta$ -ocimene, $\gamma$ -terpinene, $\alpha$ -pinene, furfuraldehyde, methylheptenone, 1,8-cineole, carvone, $\beta$ -thujone, 3-thujanol, geranyl acetate, eugenol, $\delta$ -cadinene, $\gamma$ -cadinene, dihydrocarvyl acetate  | 184-187           |
| 24.    | <i>A. parviflora</i>     | Camphor, germacrene D, germacrene B, artemisia ketone, 1,8-cineole, $\alpha$ -copaene linalool, lavandulol, santalyl acetate, lavandulyl acetate, $\beta$ -caryophyllene, spathulenol, caryophyllene oxide, cubenol, $\alpha$ -humulene, limonene, bicyclogermacrene, and $\alpha$ -cadinol   | 188-190           |
| 25.    | <i>A. myriantha</i>      | $\alpha$ -Eudesmol, $\beta$ -eudesmol, germacrene D, 1,8-cineole, $\beta$ -pinene oxide, $\delta$ -cadinene, chrysanthenone, camphor, $\beta$ -caryophyllene, $\alpha$ -humulene, caryophyllene oxide, terpinen-4-ol, and (+)- $\alpha$ -terpineol  | 191               |

(Continued)

Table 2. Continued

| S. no. | <i>Artemisia</i> species  | Essential oil components   | References |
|--------|---------------------------|--|------------|
| 26.    | <i>A. argyi</i>           | Sesquiterpenes, esters, monoterpenes, ketones, 1,8-cineole, 4-hydroxy-4-methyl-2-pentanone, borneol, thujone, $\beta$ -caryophyllene, 2,2'-bithiophene, 3,3,6,8-tetramethyl-1-tetralone and selina-6-en-4-ol, artemisia alcohol, camphor, artemisia ketone, L-borneol, absinthol, 1R- $\alpha$ -pinene, globulol, 3,3,6-trimethyl-1,5-heptadien-4-ol, isobornyl formate, spathulenol, seychellene, santolina triene, chamazulene, <i>p</i> -mentha-1,8-dien-10-ol  | 192-195    |
| 27.    | <i>A. monosperma</i>      | $\beta$ -Pinene, $\alpha$ -terpinolene, limonene, $\alpha$ -pinene, $\beta$ -maaliene, shyobunone  | 196        |
| 28.    | <i>A. sieberi</i>         | $\alpha$ -Terpineol, 1,8 cineole, $\beta$ -thujone, <i>cis</i> -sabinol, linalool, dihydrocarveol, geranyl acetate, camphor, camphene, and $\alpha$ -pinene  | 197,198    |
| 29.    | <i>A. fukudo</i>          | $\alpha$ -Thujone, $\beta$ -thujone, camphor, $\beta$ -caryophyllene and $\beta$ -elemene  | 199        |
| 30.    | <i>A. frigida</i>         | $\alpha$ -Pinene, camphene, 1,8-cineol, camphor, borneol, terpinen-4-ol, bornyl acetate, germacrene D, $\alpha$ - and $\beta$ -thujones  | 200        |
| 31.    | <i>A. argyrophylla</i>    | Yomogi alcohol, artemisia ketone, artemisia alcohol, camphor, borneol, and bornyl acetate  | 200        |
| 32.    | <i>A. imponens</i>        | Monoterpene and sesquiterpene and vulgarone B, 1,8-cineol, and camphor   | 201        |
| 33.    | <i>A. selengensis</i>     | 1,8-Cineol, ( <i>Z</i> )- $\beta$ -farnesene, $\beta$ -thujone, $\alpha$ -humulene, $\gamma$ -elemene, ( <i>E</i> )- $\beta$ -farnesene, valeranone, (2 <i>E</i> ,13 <i>Z</i> )-octadecadien-1-ol, $\alpha$ -bisabolol, bornyl acetate, germacrene D   | 202,203    |
| 34.    | <i>A. indica</i>          | $\beta$ -Caryophyllene, germacrene D, caryophyllene oxide, <i>cis</i> - $\beta$ -elemenone, and selin-11-en-4 $\alpha$ -ol   | 204        |
| 35.    | <i>A. lavandulaefolia</i> | $\beta$ -Caryophyllene, ( <i>E</i> )- $\beta$ -farnesene, and chrysanthenone   | 205        |
| 36.    | <i>A. rutifolia</i>       | 1,8-Cineol, camphor, terpenen-4-ol, and 4-isobutylphenol   | 206        |
| 37.    | <i>A. pallens</i>         | Pentanol, hexanol, ( <i>E/Z</i> )-hexenol, 2-furaldehyde/myrcene, benzyl alcohol, caprylic acid, geranyl acetone, $\alpha$ -ionone   | 207        |
| 38.    | <i>A. glauca</i>          | Anisole, methyl oleate, palmitic acid, and methyleugenol   | 208        |
| 39.    | <i>A. vestita</i> Wall    | Monoterpenes, monoterpene derivatives, sesquiterpenes, 1,8-cineol, camphor, and borneol  | 209        |
| 40.    | <i>A. mongolica</i> Fisch | Sesquiterpenes and sesquioxides, carvone, piperitone, elemol, 2-methyl-2-butene, methylene cyclopentane, $\beta$ -pinene, $\alpha$ -thujene, $\alpha$ -pinene, camphene, 1-octen-3-ol, $\beta$ -thujene, $\beta$ -pinene, $\alpha$ -phellandrene, bornylene, <i>p</i> -cymene, terpinen-1-ol, artemisia ketone, $\gamma$ -terpinene, $\beta$ -terpineol, myrtanol, $\alpha$ -terpinolene, verbenone, linalool, $\alpha$ -thujone, $\beta$ -thujone, camphor, isopulegone, isoborneol, terpinen-4-ol, $\alpha$ -terpineol, myrtenol, and <i>trans</i> - and <i>cis</i> -carveol | 210,211    |
| 41.    | <i>A. lactiflora</i> Wall | $\delta$ -3-Carene, myrcene, limonene, <i>p</i> -cymene, $\delta$ -elemene, $\alpha$ -copaene, $\beta$ -elemene, $\beta$ -caryophyllene, $\alpha$ -humulene, <i>ar</i> -curcumene, $\gamma$ -cadinene, $\delta$ -cadinene, calamenene, ( <i>E</i> )- $\beta$ -farnesene, caryophyllene oxide, $\beta$ -guaiene, lactiflorenol, spathulenol, <i>S</i> -guaiazulene, herniarin, $\alpha$ -pinene, $\beta$ -pinene, borneol, palmitic acid  | 212        |
| 42.    | <i>A. subdigitata</i>     | $\alpha$ -Thujone, $\alpha$ -pinene, camphene, sabinene, $\beta$ -pinene, myrcene, $\delta$ -3-carene, $\beta$ -ocimene, <i>p</i> -isopropylphenol, limonene, $\gamma$ -terpinene, terpinen-4-ol, estragole, geraniol, methyl eugenol  | 213        |

*Artemisia japonica* Thunb. is distributed throughout continental Asia and Japan.<sup>37</sup> In northern Pakistan, the leaf extract is used to treat malaria and a paste of leaves is used externally on skin diseases.<sup>35</sup> *Artemisia judaica* L. is a perennial fragrant shrub that grows widely in the deserts and on the Sinai Peninsula in Egypt, and is a very common anthelmintic drug in most North African and Middle-Eastern countries where it is known by the Arabic name of "shih."<sup>2,22</sup>


The genus *Artemisia* is known to contain many bioactive compounds; artemisinin exerts not only antimalarial activity but also profound cytotoxicity against tumor cells<sup>6,42</sup> and arglabin is employed for treating certain types of cancer in the former Soviet Union.<sup>43</sup> Over the past decade *Artemisia* species have been used traditionally in various populations; thus, *A. keiskeana* Miq. has been used as a traditional Chinese drug for the treatment of gynecopathy, amenorrhea, bruise, and rheumatic disease.<sup>6,44</sup>

The inhabitants of northeastern Mexico use an infusion of leaves from *A. ludoviciana* Nutt. as an antidiarrheal remedy,<sup>2</sup> while several Native American tribes of North America have used infusions for coughs, sore throats, and colds.<sup>39</sup> *Artemisia nilagirica* (C.B. Clarke) Pamp., commonly called "Indian wormwood," is widely found in the hilly areas of India, where it is used as insecticide.<sup>45</sup> A paste from the leaves of *A. nilagirica* is used externally to treat cuts and wounds and the leaves chewed to treat oral ulcers.<sup>35</sup>

*Artemisia princeps* Willd. ("Japanese mugwort" or "yomogi") is the best-known *Artemisia* in Japan, where it is a fundamental ingredient of the Japanese confection "kusa-mochi." This plant has also been used in traditional Asian medicine for the treatment of inflammation, diarrhea, and many circulatory disorders.<sup>2</sup>

*Artemisia rubripes* Nakai has been used as a traditional Korean medicine for stomach ache, vomiting, diarrhea, and as a





**Figure 1.** Some nonvolatile constituents of *Artemisia* spp.

hemostatic agent.<sup>46</sup> *Artemisia rutifolia* Stephan ex Spreng. is distributed in Afghanistan, China, India, Kazakhstan, Kyrgyzstan, Mongolia, Nepal, Pakistan, Russian Federation, and Tajikistan. A tea from the herb is taken to treat asthma, as an anti-inflammatory and diuretic; the fresh herb is used as an analgesic for toothache; a decoction of the herb is gargled for treatment of angina, stomach problems, and heart problems.<sup>14,47</sup>

*Artemisia scoparia* Waldst. & Kit. (“red stem worm wood”) is a faintly scented annual herb which is widespread and common throughout the world, particularly in southwest Asia and central Europe. The success of *A. scoparia* may be attributed to the presence of phytotoxins, the volatile essential oils, in addition to other nonvolatile secondary products. It has been established that aerial parts of *A. scoparia* yield a volatile essential oil that has medicinal value. It possesses insecticidal, antibacterial, anticholesterolemic, antipyretic, antiseptic, cholagogue, diuretic, purgative, and vasodilatory activities and is also used for the treatment of gall bladder inflammation, hepatitis, jaundice, malaria, and diabetes.<sup>2,35,48</sup>

*Artemisia spicigera* K. Koch, named locally as “yavs, an,” is widespread in Central and Eastern Anatolia in Turkey, at an altitude between 1000 and 2500 m.<sup>49</sup>

*Artemisia tridentata* Nutt. “big sagebrush” is one of the most widely distributed and ecologically important shrub species in Western North America. This species serves as a critical habitat and food resource for many animals and invertebrates.<sup>32</sup> Several North American native tribes have used infusions of *A. tridentata* to treat bronchitis and pneumonia.<sup>39</sup>

*Artemisia vestita* Wall ex Besser has been utilized for the treatment of fungal infections such as tinea, tympanitis, and thrush.<sup>2,6,23</sup>

*Artemisia vulgaris* L., commonly known as “mugwort,” is a perennial weed growing wild native in temperate and cold-temperature zones of the world such as in Asia, Europe, and North America.<sup>50,51</sup> The plant is widely used in the Philippines, where it is locally known as “herbaka,” for its antihypertensive actions. It has also been suggested to have

other medicinal activities such as anti-inflammatory, anti-spasmodic, carminative, and anthelmintic properties, and has been used in the treatment of painful menstruation (dysmenorrhea) and in the induction of labor or miscarriage.<sup>50</sup> *Artemisia vulgaris* has been known not only as an edible plant but also as a folk medicine resource. Mugwort is used to flavor tea and rice dishes in Asia and as a culinary herb for poultry and pork in Western cultures.<sup>51</sup> In Oriental medicine, mugwort has been employed as an analgesic agent and in conjunction with acupuncture therapy.<sup>51,52</sup> Considered an emmenagogue, an inducing agent of menstrual flow, mugwort has been traditionally employed to bring about regular menses in cases of amenorrhea or menorrhagia.<sup>51</sup>

*Artemisia mongolica* (Fisch. ex Besser) Nakai has been used as a folk medicine for generations to cure inflammations and colds in Northwest China.<sup>53</sup>

*Artemisia pontica* L. is well known in Bulgarian folk medicine as a sedative and an appetizer.<sup>54</sup>


In the western Himalayas of northern Pakistan, *A. roxburghiana* Besser extract is used to treat fever, malaria, and intestinal worms. In Uttarakhand, India, *A. roxburghiana* is used in ethnoveterinary medicine to treat eye diseases, wounds, cuts, and external parasites.<sup>35</sup>

*Artemisia herba-alba* Asso (syn. *Artemisia maritima* L., *Artemisia brevifolia* Wall.) is used in the traditional medicine of the Northern Badia region of Jordan, in the form of a decoction, against fever and menstrual and nervous problems.<sup>55</sup> *Artemisia herba-alba*, known also as desert wormwood (known in Arabic as “shih,” and French as “armoiseblanche”),<sup>56</sup> has been used in folk medicine by many cultures since ancient times and in Moroccan folk medicine to treat arterial hypertension and/or diabetes.<sup>57-59</sup> *Artemisia herba-alba* is widely distributed in North Africa and used traditionally by the Egyptians as a vermifuge in addition to its other medical and veterinary uses.<sup>60</sup> Herbal tea from this species has been used as analgesic, antibacterial, antispasmodic, and hemostatic agents.<sup>61</sup> During an ethnopharmacological survey carried out among the Bedouins of the Negev Desert, it was found that *A. herba-alba* relieved stomach disorders.<sup>60</sup> This plant is also suggested to be important as a fodder for sheep and for livestock in the plateau regions of Algeria where it grows abundantly.<sup>62</sup>

*Artemisia sieberi* Besser is a famous medicinal plant in Middle East traditional medicine as an anthelmintic. In external use, the flowering shoots and leaves were boiled in normal saline and the extracted solution was used for treatment of gangrenous ulcers, infectious ulcers, and inflammations. *Artemisia sieberi* is used as fodder for sheep and it is believed that it can increase weight and fleece of sheep. It was used as carminative, to relieve inflammation and abscesses and to prevent leprosy.

## Bioactive Compounds From *Artemisia*

Bioactive compounds are experiencing a growing interest in wide range of applications: geo-medicine, plant science,



**Figure 2.** Structure of some volatile constituents of *Artemisia* spp.

modern pharmacology, agrochemicals, cosmetics, food industry, nano-bio-science, and so on.<sup>63-65</sup>

Bioactive compounds in plants are classified according to different criteria. A presentation based on clinical function—their pharmacological or toxicological effects—is relevant for the clinician, pharmacist, or toxicologist. An approach based on biological effects is complicated by the fact that the clinical outcome is not exclusively connected to biochemically closely related compounds; even biochemically different molecules might produce similar clinical effects. A botanical categorization based on families and genera of the plants producing the bioactive compounds might also be relevant, as closely related plant species most often produce the same or chemically similar bioactive compounds. However, there are also ranges of examples that species even genetically less related produce similar secondary compounds. The main focus are the bioactive chemical compounds; therefore, it is useful to categorize them according to biochemical pathways and chemical classes.<sup>66-68</sup>

*Artemisia* species represent rich sources of various types of biologically active compounds accountable for numerous pharmacological activities. Differences in qualitative and quantitative composition of bioactive compounds might be correlated with environmental conditions, species variation, geographic, climatic, and genetic conditions, plant age, soil, phase of vegetation, anatomical part of plant, harvesting season, and method of harvesting.<sup>69-72</sup>

Tables 1 and 2 present detailed list of the chemical constituents and essential oil components of various *Artemisia* sp., respectively (Figures 1 and 2).

## Conclusion

*Artemisia* species are widely used in traditional medicine all over the world with different and well-known therapeutic applications. They exhibit anti-inflammatory, antitumor, antioxidant,

antispasmodic, antimicrobial, insecticidal, antimalarial, antifungal, and antioxidant activities. These diverse biological activities are manifested by different compounds whose main components are essential oils and polyphenols. *Artemisia* holds a great potential for human health and its therapeutic effects should be more strictly and intensively analyzed. Preclinical and clinical research needs to be done on the use of these plants and further indepth investigations are urgently necessary to study all bioactive compounds and their biomolecular mechanisms at the cellular and tissue levels.

### Authors' Note

Sergey Plygun is now affiliated with Laboratory of Biocontrol and Antimicrobial Resistance, Orel State University named after I.S. Turgenyev, Orel, Russia and European Society of Clinical Microbiology and Infectious Diseases, Basel, Switzerland. Javad Sharifi-Rad is now affiliated with Zabol Medicinal Plants Research Center, Zabol University of Medical Sciences, Zabol, Iran.

### Acknowledgments

The authors are very thankful to all the authors whose work has been cited in this paper.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

### References

- Bishop JF, Matthews JP, Young GA, et al. A randomized study of high-dose cytarabine in induction in acute myeloid leukemia. *Blood*. 1996;87(5):1710-1717.
- Abad MJ, Bedoya LM, Apaza L, Bermejo P. The *Artemisia* L. genus: a review of bioactive essential oils. *Molecules*. 2012;17(3):2542-2566.
- Bora KS, Sharma A. The genus *Artemisia*: a comprehensive review. *Pharm Biol*. 2011;49(1):101-109.
- Teixeira da Silva JA. Mining the essential oils of the Anthemideae. *Afr J Biotechnology*. 2004;3(12):706-720.
- Willcox M. *Artemisia* species: From traditional medicines to modern antimalarials--and back again. *J Altern Complement Med*. 2009;15(2):101-109.
- Mohamed AH, El-Sayed MA, Hegazy ME, Helaly SE, Esmail AM, Mohamed NS. Chemical constituents and biological activities of *Artemisia herba-alba*. *Rec Nat Products*. 2010;4:1-25.
- Mohsen H, Ferchichi A. Study of genetic polymorphism of *Artemisia herba-alba* from Tunisia using ISSR markers. *Afr J Biotechnol*. 2008;7(1):44-50.
- Ferreira J, Janick J. 2009. *Annual Wormwood (Artemisia annua L)*. [www.hort.purdue.edu/newcrop/cropfactsheets/artemisia.pdf](http://www.hort.purdue.edu/newcrop/cropfactsheets/artemisia.pdf)
- Wright CW. *Artemisia. Medicinal and Aromatic Plants - Industrial Profiles*. London: CRC Press; 2003:1-344.
- Salehi B, Sharopov F, Martorell M, et al. Phytochemicals in *Helicobacter pylori* infections: what are we doing now? *Int J Mol Sci*. 2018;19(8):2361.
- Sharifi-Rad M, Nazaruk J, Polito L, et al. *Matricaria* genus as a source of antimicrobial agents: From farm to pharmacy and food applications. *Microbiol Res*. 2018;215:76-88.
- Sharifi-Rad J, Sureda A, Tenore G, et al. Biological activities of essential oils: from plant chemocology to traditional healing systems. *Molecules*. 2017;22(1):70.
- Sharopov FS, Sulaimonova VA, Setzer WN. Composition of the essential oil of *Artemisia absinthium* from Tajikistan. *Rec Nat Products*. 2012;6:127-134.
- Sharopov FS, Zhang H, Wink M, Setzer WN. Aromatic medicinal plants from Tajikistan (Central Asia). *Medicines*. 2015;2(1):28-46.
- Lachenmeier DW. Wormwood (*Artemisia absinthium* L.)-a curious plant with both neurotoxic and neuroprotective properties? *J Ethnopharmacol*. 2010;131(1):224-227.
- Çubukçu B, Bray DH, Warhurst DC, Meriçli AH, Özhatay N, Sarıyay G. In vitro antimalarial activity of crude extracts and compounds from *Artemisia abrotanum* L. *Phytother Res*. 1990;4(5):203-204.
- Mahboubi M. *Artemisia sieberi* Besser essential oil and treatment of fungal infections. *Biomed Pharmacother*. 2017;89:1422-1430.
- Chhetri BK, Ali NAA, Setzer WN. A survey of chemical compositions and biological activities of Yemeni aromatic medicinal plants. *Medicines*. 2015;2(2):67-92.
- Tadesse M. *Flora of Ethiopia and Eritrea: Asteraceae (Compositae)*. Addis Ababa University, Ethiopia: National Herbarium Biology Department; 2005.
- Chhetri BK, Al-Sokari SS, Setzer WN, Awadh Ali NA. Essential oil composition of *Artemisia abyssinica* from three habitats in Yemen. *Am J Essent Oil Nat Prod*. 2015;2(3):28-30.
- Bergendorff O, Sterner O. Spasmolytic flavonols from *Artemisia abrotanum*. *Planta Med*. 1995;61(4):370-371.
- van Wyk B-E. A broad review of commercially important southern African medicinal plants. *J Ethnopharmacol*. 2008;119(3):342-355.
- Klayman DL. *Artemisia annua*: from weed to respectable antimalarial plant. In: Kinghorn AD, Balandrin MF, eds. *Human Medicinal Agents from Plants. ACS symposium series (USA)*; 1993:242-255.
- Hutchings A. *Zulu Medicinal Plants: An Inventory*. South Africa: University of Natal Press; 1996.
- Watt JM, Breyer-Brandwijk MG. *The medicinal and poisonous plants of southern and eastern Africa: being an account of their medicinal and other uses, chemical composition, pharmacological effects and toxicology in man and animal*. E. & S. USA: Livingstone; 1962.
- Hsu E. The history of Qing Hao in the Chinese materia MEDICA. *Trans R Soc Trop Med Hyg*. 2006;100(6):505-508.
- Pinheiro LCS, Feitosa LM, Silveira FFDA, Boechat N. Current antimalarial therapies and advances in the development of semi-synthetic artemisinin derivatives. *An Acad Bras Ciênc*. 2018;90(1 suppl 2):1251-1271.

28. Sadiq A, Hayat MQ, Ashraf M. Ethnopharmacology of *Artemisia annua* L.: A Review. In: Aftab T, Ferreira JFS, Khan MMA, Naeem M, eds. *Artemisia annua - Pharmacology and Biotechnology*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2014:9-25.
29. Salehi B, Kumar N, Şener B, et al. Medicinal plants used in the treatment of human immunodeficiency virus. *Int J Mol Sci*. 2018;19(5):1459.
30. Ballero M, Poli F, Sacchetti G, Loi MC. Ethnobotanical research in the territory of Fluminimaggiore (south-western Sardinia). *Fitoterapia*. 2001;72(7):788-801.
31. Edward F. *Contribution une étude ethnobotanique de la flore tunisienne*. Imprimerie Officielle de la République Tunisienne: Tunisia; 1983.
32. Moerman DE. An analysis of the food plants and drug plants of native North America. *J Ethnopharmacol*. 1996;52(1):1-22.
33. Setzer WN, Vogler B, Schmidt JM, Leahy JG, Rives R. Antimicrobial activity of *Artemisia douglasiana* leaf essential oil. *Fitoterapia*. 2004;75(2):192-200.
34. Espinar LA, Bonzani N. El mático de la Región de Cuyo (Argentina). *Acta Farm Bonaer*. 1992;11:139-145.
35. Joshi R, Satyal P, Setzer W. Himalayan aromatic medicinal plants: a review of their Ethnopharmacology, volatile Phytochemistry, and biological activities. *Medicines*. 2016;3(1):6.
36. Obolskiy D, Pischel I, Feistel B, Glotov N, Heinrich M. *Artemisia dracunculus* L. (tarragon): a critical review of its traditional use, chemical composition, pharmacology, and safety. *J Agric Food Chem*. 2011;59(21):11367-11384.
37. Wu ZY, Raven PH. *Flora of China*. Beijing & St. Louis: Science Press & Missouri Botanical Garden Press; 2011.
38. Satyal P, Paudel P, Kafle A, et al. Bioactivities of volatile components from Nepalese *Artemisia* species. *Nat Prod Commun*. 2012;7(12):1651-1658.
39. Moerman DE. *Native American Ethnobotany, 1st ed*. Portland: USA: Timber Press, Incorporated; 1998.
40. Lee YS, Park JY, Kim J-K, et al. Complete chloroplast genome sequence of *Artemisia fukudo* Makino (Asteraceae). *Mitochondrial DNA Part B*. 2016;1(1):376-377.
41. Khanahmadi M, Rezazadeh S. Review on Iranian medicinal plants with antioxidant properties. *J Med Plants*. 2010;9(35):19-32.
42. Efferth T. Willmar Schwabe Award 2006: antiplasmodial and antitumor activity of artemisinin--from bench to bedside. *Planta Med*. 2007;73(4):299-309.
43. Wong HF, Brown GD. Germacranolides from *Artemisia myriantha* and their conformation. *Phytochemistry*. 2002;59(5):529-536.
44. Kwak JH, Jang WY, Zee OP, Lee KR. Artekeiskeanin A: a new coumarin-monoterpene ether from *Artemisia keiskeana*. *Planta Med*. 1997;63(5):474-476.
45. Bhattacharjee SK. *Handbook of Medicinal Plants*. Jaipur, India: Aavishkar Publishers; 2000.
46. Lee KH, Min YD, Choi SZ, et al. A new sesquiterpene lactone from *Artemisia rubripes* nakai. *Arch Pharm Res*. 2004;27(10):1016-1019.
47. Sharopov FS, Setzer WN. Thujone-rich essential oils of *Artemisia rutifolia* Stephan ex Spreng. growing wild in Tajikistan. *J Essent Oil Bear Pl*. 2011;14(2):136-139.
48. Sharopov FS, Setzer WN. The essential oil of *Artemisia scoparia* from Tajikistan is dominated by phenyldiacetylenes. *Nat Prod Commun*. 2011;6(1):119-122.
49. Kordali S, Kotan R, Mavi A, Cakir A, Ala A, Yildirim A. Determination of the chemical composition and antioxidant activity of the essential oil of *Artemisia dracunculus* and of the antifungal and antibacterial activities of Turkish *Artemisia absinthium*, *A. dracunculus*, *Artemisia santonicum*, and *Artemisia spicigera* essential oils. *J Agric Food Chem*. 2005;53(24):9452-9458.
50. Quisumbing E. *Medicinal Plants of the Philippines*. Manila, Philippines: Bureau of Printing; 1978.
51. Lee S-J, Chung H-Y, Maier CG-A, Wood AR, Dixon RA, Mabry TJ. Estrogenic flavonoids from *Artemisia vulgaris* L. *J Agric Food Chem*. 1998;46(8):3325-3329.
52. Yoshikawa M, Shimada H, Matsuda H, Yamahara J, Murakami N. Bioactive constituents of Chinese natural medicines. I. new sesquiterpene ketones with vasorelaxant effect from Chinese moxa, the processed leaves of *Artemisia argyi* Levl. et Vant.: moxartenone and moxartenolide. *Chem Pharm Bull*. 1996;44(9):1656-1662.
53. Hu J, Zhu Q, Bai S, Jia Z. New eudesmane sesquiterpene and other constituents from *Artemisia mongolica*. *Planta Med*. 1996;62(5):477-478.
54. Todorova MN, Tsankova ET, Trendafilova AB, Gussev CV. Sesquiterpene lactones with the uncommon rotundane skeleton from *Artemisia pontica* L. *Phytochemistry*. 1996;41(2):553-556.
55. Alzweiri M, Sarhan AA, Mansi K, Hudaib M, Aburjai T. Ethnopharmacological survey of medicinal herbs in Jordan, the Northern badia region. *J Ethnopharmacol*. 2011;137(1):27-35.
56. Segal R, Feuerstein I, Danin A. Chemotypes of *Artemisia herba-alba* in Israel based on their sesquiterpene lactone and essential oil constitution. *Biochem Syst Ecol*. 1987;15(4):411-416.
57. Ziyayt A, Legssyer A, Mekhfi H, Dassouli A, Serhrouchni M, Benjelloun W. Phytotherapy of hypertension and diabetes in Oriental Morocco. *J Ethnopharmacol*. 1997;58(1):45-54.
58. Tahraoui A, El-Hilaly J, Israili ZH, Lyoussi B. Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in south-eastern Morocco (Errachidia Province). *J Ethnopharmacol*. 2007;110(1):105-117.
59. Zeggwagh N-A, Farid O, Michel JB, Eddouks M. Cardiovascular effect of *Artemisia herba alba* aqueous extract in spontaneously hypertensive rats. *Methods Find Exp Clin Pharmacol*. 2008;30(5):375-381.
60. Seddiek SA, Ali MM, Khater HF, El-Shorbagy MM. Anthelmintic activity of the white wormwood, *Artemisia herba-alba* against *Heterakis gallinarum* infecting turkey poults. *J Med Plant Res*. 2011;5:3946-3957.
61. Laid M, Hegazy MEF, Ahmed AA, Ali K, Belkacemi D, Ohta S. Sesquiterpene lactones from Algerian *Artemisia herba-alba*. *Phytochem Lett*. 2008;1(2):85-88.

62. Belhattab R, Boudjouref M, Barroso JG, Pedro LP, Figueirido AC. Essential oil composition from *Artemisia campestris* grown in Algeria. *Adv Environ Biol.* 2011;5:429-432.
63. Sharifi-Rad M, Roberts TH, Matthews KR, et al. Ethnobotany of the genus *Taraxacum*-Phytochemicals and antimicrobial activity. *Phytother Res.* 2018;32(11):2131-2145.
64. Salehi B, Valussi M, Jugran AK, et al. *Nepeta* species: from farm to food applications and phytotherapy. *Trends Food Sci Technol.* 2018;80:104-122.
65. Sharifi-Rad M, Fokou PVT, Sharopov F, et al. Antiulcer agents: from plant extracts to phytochemicals in healing promotion. *Molecules.* 2018;23(7):1751.
66. Mishra AP, Sharifi-Rad M, Shariati MA, et al. Bioactive compounds and health benefits of edible *Rumex* species-A review. *Cell Mol Biol.* 2018;64(8):27-34.
67. Mishra AP, Saklani S, Salehi B, et al. *Satyrium nepalense*, a high altitude medicinal orchid of Indian Himalayan region: chemical profile and biological activities of tuber extracts. *Cell Mol Biol.* 2018;64(8):35-43.
68. Sharifi-Rad M, Ozcelik B, Altun G, et al. *Salvia* spp. plants-from farm to food applications and phytopharmacotherapy. *Trends Food Sci Technol.* 2018;80(9):242-263.
69. Burbott AJ, Loomis WD. Effects of light and temperature on the monoterpenes of peppermint. *Plant Physiol.* 1967;42(1):20-28.
70. Gudaitytė O, Venskutonis PR. Chemotypes of *Achillea millefolium* transferred from 14 different locations in Lithuania to the controlled environment. *Biochem Syst Ecol.* 2007;35(9):582-592.
71. Kokkini S, Karousou R, Vokou D. Pattern of geographic variations of *Origanum vulgare* trichomes and essential oil content in Greece. *Biochem Syst Ecol.* 1994;22(5):517-528.
72. Tétényi P. Chemical variation (Chemodifferentiation) in medicinal and aromatic plants. *International Conference on Medicinal and Aromatic Plants. Possibilities and Limitations of Medicinal and Aromatic Plant 576*; 2002.
73. Zhang X, Zhao Y, Guo L, et al. Differences in chemical constituents of *Artemisia annua* L from different geographical regions in China. *PLoS One.* 2017;12(9):e0183047.
74. Qingjuan LI, Chen W, Fan J, et al. Study on the chemical constituents of *Artemisia myriantha* Wall.Ex Bess. *Chin J Mod Appl Pharm.* 2014;31:706-710.
75. Anshul N, Bhakuni RS, Gaur R, Singh D. Isomeric flavonoids of *Artemisia annua* (Asterales: Asteraceae) as insect growth inhibitors against *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Fla Entomol.* 2013;96(3):897-903.
76. Shen L, Shi D-H, Song Y-C, Tan R-X. Chemical constituents of liquid culture of endophyte IFB-E012 in *Artemisia annua*. *Chin J Nat Med.* 2009;7(5):354-356.
77. Jing C, Yubo Z, Xin Z, Li H, Wei S, Jinhui W. Chemical constituents of young leaves of *Artemisia annua*. *J Sheny Pharm Univ.* 2008;25:866-870.
78. Al-Wahaibi LHN, Mahmood A, Khan M, Alkhatlan HZ. Comparative study on the essential oils of *Artemisia judaica* and *A. herba-alba* from Saudi Arabia. *Arab J Chem.* 2018.
79. Pandey BP, Thapa R, Upreti A. Chemical composition, antioxidant and antibacterial activities of essential oil and methanol extract of *Artemisia vulgaris* and *Gaultheria fragrantissima* collected from Nepal. *Asian Pac J Trop Med.* 2017;10(10):952-959.
80. Van Nguyen Thien T, Tran LTK, Nhu NTT, et al. A new eudesmane-type sesquiterpene from the leaves of *Artemisia vulgaris*. *Chem Nat Compd.* 2018;54(1):66-68.
81. Gairola S, Maithani M, Gupta V, Bansal P, Ghaiye P. Pharmacological potential and chemical constituents of *Artemisia vulgaris*. *J Pharm Res & Clinical Practice.* 2011;1:82-87.
82. Wu T, He F, Ma QL, Chen J, Aisa HA. Chemical constituents of *Artemisia rupestris*. *Chem Nat Compd.* 2017;53(5):991-993.
83. Suzhang Z, Wei Y, Zhengyi G. Study on the chemical constituents of *Artemisia scoparia*. *J Xinjiang Med Univ.* 2016;39:408-410.
84. Song W-X, Ji T-F, Si Y-K, Su Y-L, , YK S, YL S. Studies on chemical constituents in herb from *Artemisia rupestris*. *Zhongguo Zhong Yao Za Zhi.* 2006;31(21):1790-1792.
85. Yu F, Qian H, Zhang J, Sun J, Ma Z. Simultaneous quantification of eight organic acid components in *Artemisia capillaris* Thunb (Yinchen) extract using high-performance liquid chromatography coupled with diode array detection and high-resolution mass spectrometry. *J Food Drug Anal.* 2018;26(2):788-795.
86. Geng C-A, Huang X-Y, Chen X-L, et al. Three new anti-HBV active constituents from the traditional Chinese herb of Yin-Chen (*Artemisia scoparia*). *J Ethnopharmacol.* 2015;176:109-117.
87. Li-Hong W, Yan X, Yu-Xi Z, Xiao-Ying Y, Yang YJS. Isolation and identification of chemical constituents from *Artemisia capillaris*. *J China pharm.* 2011;22:1020-1022.
88. Hongyu M, Yi S, Ali L, Gang C, Honghua W, Lake P. Isolation and identification of chemical components from *Artemisia capillaris*. *Chin J Med Chem.* 2010;20:61-63+69.
89. Zhiwei W, Xiaojie T, Tingting M, Xiaohui C, Kaishun B. Isolation and identification of chemical components of Yinchen. *J Shenyang Pharm Univ.* 2008;25:781-784.
90. Pandey AK, Singh P. The genus *Artemisia*: A 2012–2017 literature review on chemical composition, antimicrobial, insecticidal and antioxidant activities of essential oils. *Medicines.* 2017;4(3):68.
91. Zhang W, Zhao D-bao, Li M-jing, Liu X-hua, Wang H-qing, MJ L, Wang HQ. Studies on flavonoid constituents from herbs of *Artemisia ordosica* II. *Zhongguo Zhong Yao Za Zhi.* 2006;31(23):1959-1961.
92. Govindarajan M, Benelli G. *Artemisia absinthium*-borne compounds as novel larvicides: effectiveness against six mosquito vectors and acute toxicity on non-target aquatic organisms. *Parasitol Res.* 2016;115(12):4649-4661.
93. Lee H-G, Kim H, Oh W-K, et al. Tetramethoxy hydroxyflavone p7F downregulates inflammatory mediators via the inhibition of nuclear factor kappaB. *Ann N Y Acad Sci.* 2004;1030:555-568.
94. Bora KS, Sharma A. Phytochemical and pharmacological potential of *Artemisia absinthium* Linn. and *Artemisia asiatica* Nakai : A Review. *J Pharm Res.* 2016;3:325-328.
95. Zhang J, Li L, Liu X, Wang Y, Zhao D. Study on chemical constituents of *Artemisia sphaerocephala*. *Zhongguo Zhong Yao Za Zhi.* 2012;37(2):238-242.

96. Weiyu W, Tongshi W, Hui L, Dayuan Z, Jixiang J, Yuhui L. Study on chemical constituents of sand-fixing plant *Artemisia sphaerocephala* [I]. *J Northwest Normal Univ.* 1988;18:60-62.
97. Mojarrab M, Saremi G, Emami SA. Evaluation of antioxidant activity and identification of main compounds of various extracts of *Artemisia turanica* aerial parts. *Res J Pharm.* 2017;4:36.
98. Safari S, Taherkhani M. Extraction and identification of flavon from *Artemisia turanica* Krasch the extract which has been collected from Esfarayen, Khorasan province. *Eco-phytochem J Med Pl.* 2018;6:44-55.
99. Al Jahid A, Essabaq S, Elamrani A, Blaghen M, Jamal Eddine J. Chemical composition, antimicrobial and antioxidant activities of the essential oil and the hydro-alcoholic extract of *Artemisia campestris* L. leaves from southeastern Morocco. *J Bio Active Prod Nat.* 2016;6(5-6):393-405.
100. Zeng W, Liang H. Flavonoids from *Artemisia gmelinii* Web. ex Stechm. *J Chin Pharm Sci.* 2014;23(7).
101. He-Xang D, Wen-Yan L, Yi-Sheng Y, Qi-Zhen C, Xu-Ping L, Hong-Gang H. Chemical constituents in ethyl acetate fraction of *Artemisia selengensis*. *Chin J Integr Med.* 2015;46:1441-1444.
102. Kim AR, Ko HJ, Chowdhury MA, Chang Y-S, Woo E-R. Chemical constituents on the aerial parts of *Artemisia selengensis* and their IL-6 inhibitory activity. *Arch Pharm Res.* 2015;38(6):1059-1065.
103. Lili C, Zu-Min Q, Zhen-Zhong H, Wen-You C. Study on the chemical constituents of volatile oil from the shoots of *Artemisia selengensis* by GC-MS. *J Nanchang Univ.* 2008;30:212-214.
104. Jian Z, Lingyi K. Study on chemical constituents of *Artemisia selengensis* leaves. *Chin J Pharm.* 2005;40:1778-1780.
105. Lin M, Hailing Z, Youming G, Li H, Qi D. Study on flavonoids in *Artemisia selengensis*. *Tradit Chin Drug Res Clin Pharmacol.* 2012;23:555-557.
106. Qun-Hui L, Nai-Li W, Hong-Wei L, Ming F, Ai-Shi D, Xin-Sheng Y. Chemical constituents from *Artemisia scoparia*. *J Shenyang Pharm Univ.* 2006;23:492-494.
107. Lin S, Xiao Y-qing, Zhang Q-wei, Zhang N-ning. Studies on chemical constituents in bud of *Artemisia scoparia* (II). *Zhongguo Zhong Yao Za Zhi.* 2004;29(2):152-154.
108. Lin S, Xiao Y-qing, Zhang Q-wei, Shi J-gong, Wang Z-min. Studies on chemical constituents in bud of *Artemisia scoparia* (III). *Zhongguo Zhong Yao Za Zhi.* 2004;29(5):429-431.
109. Wei X, Jingwei L, Jing L, Min W, Xiuli W, Chunhua Y. Study on the chemical constituents of *Artemisia scoparia*. *J China Pharm Univ.* 2004;35:401-403.
110. Wei X, Jing L, Zhiming Z, Xiuli W. Study on *Artemisia alkeyne* and Flavonoids II. *Chin J Nat Med.* 2005;3:86-89.
111. Zhang QW, Zhang YX, Zhang Y, Xiao YQ, Wang ZM. Studies on chemical constituents in buds of *Artemisia scoparia*. *Zhongguo Zhong Yao Za Zhi.* 2002;27:202-204.
112. Wang QH, XL W, Wang JH. Chemical constituents of *Artemisia frigida* (II). *Chin Tradit Herbal Drug.* 2011;42:1075-1078.
113. Lin F-D, Luo D-W, Ye J, Xiao M-T. Chemical constituents of *Artemisia lactiflora* (II). *Zhongguo Zhong Yao Za Zhi.* 2014;39(13):2531-2535.
114. Xiao MT, Luo DW, Zan K, Ye J, PF T. Chemical constituents from the aerial parts of *Artemisia lactiflora* (III). *J Chin Pharm.* 2015;50:209-212.
115. Mei-Tian X, Jing Y, Ben-Bo H, Ke Z, Peng-Fei T. Study on chemical constituents of *Artemisia lactiflora*. *Chin Pharm J.* 2011;46:414-417.
116. Zeng Y-T, Jiang J-M, Lao H-Y, Guo J-W, Lun Y-N, Yang M. Antitumor and apoptotic activities of the chemical constituents from the ethyl acetate extract of *Artemisia indica*. *Mol Med Rep.* 2015;11(3):2234-2240.
117. Weihong L, Wei P, Jishan F, Bin L. Study on chemical constituents of *Artemisia scoparia*. *China Pharm.* 2014;25:253-255.
118. RL X, Shi Y. Chemical constituents from *Artemisiae Anomalae* Herba. *Chin Tradit Herbal Drug.* 2014;45:1521-1525.
119. Tong-shu X, Qiong W, Li-long J, Jian-qin J, You-bin L. Chemical constituents of *Artemisia anomala*. *Chin Tradit Herbal Drug.* 2013;44:515-518.
120. Zan K, Chen X-Q, TU P-F, . A new 1, 10-secoguaianolide from the aerial parts of *Artemisia anomala*. *Chin J Nat Med.* 2012;10(5):358-362.
121. Jing W, Haiming S, Zuo Z, et al. Chemical constituents in *Artemisia anomala*. *Chin Tradit Herbal Drug.* 2010;41:870-873.
122. Furaio T, Lin Z, Kui T, Wenming Z. Study on the chemical constituents of nan Liu Sinuo. *Chin J Med Chem.* 2008;18:362-365.
123. Wang S, Tu P. Anti-neuroinflammatory constituents from *Artemisia argyi*. *J Chin Pharm Sci.* 2013;22:377-380.
124. Weiguang R, Sensen L, Wentao L, Linfang H. UPLC Q-TOF/MS method to study the active site of EGFR kinase. *Huaxi Pharmaceutical Journal.* 2013;28:604-606.
125. Ji S, Lu G, Meng D, Li N, Li X. Chemical constituents from the Folium *Artemisiae Argyi* (II). *J Shenyang Pharm Univ.* 2010;27(548-550):566.
126. Cui F-X, Zhang C, Jiang Y, Tu P-F, . Chemical constituents from ethyl acetate extract of *Artemisia rupestris*. *Zhongguo Zhong Yao Za Zhi.* 2013;38(11):1757-1759.
127. Ji T-fei, Yang J-bo, Song W-xia, Wang A-guo, Su Y-lun, Yuan L. Studies on chemical constituents of *Artemisia rupestris* (II). *Zhongguo Zhong Yao Za Zhi.* 2007;32(12):1187-1189.
128. Chong L, Jia-Xiu D, Li-Li J. Chemical constituents of petroleum ether extract from the *Artemisia sacrorum* Ledeb. *J Med Sci Yanbian Univ.* 2013;36:27-28.
129. Lijuan D, Dang D. Extraction and separation of a liver-protecting active ingredient from *Artemisia scoparia*. *Guangdong Trace Elem Sci.* 2008;15:22-23.
130. Lone SH, Bhat KA, Naseer S, Rather RA, Khuroo MA, Tasduq SA. Isolation, cytotoxicity evaluation and HPLC-quantification of the chemical constituents from *Artemisia amygdalina* Decne. *J Chromatogr B.* 2013;940:135-141.
131. Cha J-D, Jeong M-R, Choi H-J, et al. Chemical composition and antimicrobial activity of the essential oil of *Artemisia lavandulaefolia*. *Planta Med.* 2005;71(6):575-577.
132. Wang X-qin, Zhou C-jiang, Zhang N, Wu G, Li M-hui. Studies on the chemical constituents of *Artemisia lavandulaefolia*. *Zhong Yao Cai.* 2011;34(2):234-236.

133. Kiani BH, Ullah N, Haq I-ul, Mirza B. Transgenic *Artemisia dubia* WALL showed altered phytochemistry and pharmacology. *Arab J Chem*. 2015.
134. Hao-Yang Z, Ablajan K, Yan L, Zhu-Wei Y, Yang P. Chemical constituents from aerial parts of *Artemisia dracunculus* var. *turkestanica*. *Chin Tradit Herbal Drug*. 2013;44:1096-1100.
135. Yazdan Parast R, Alavi HR, Bazarganian A. Two new compounds from *Artemisia dracunculus* L. *Daru J Pharm Sci*. 2000;8:42-44.
136. Lee YK, Hong EY, Whang WK. Inhibitory effect of chemical constituents isolated from *Artemisia inuyomogi* on Polyol pathway and simultaneous quantification of major bioactive compounds. *Biomed Res Int*. 2017;12(4):1-12.
137. Yan D, Chun L, Choi EM, JC R, Kim YH. Chemical constituents from *Artemisia inuyomogi* increase the function of osteoblastic MC3T3-E1 cells. *Nat Prod Sci*. 2009;15:192-197.
138. Wang Y, Yin J, Qiao Y, Zhang H, Lu X. Studies on antioxidant activity and chemical constituents of *Artemisia halodendron*. *Asian J Tradit Med*. 2007;2:30-33.
139. Zuo X, Jing L, Jingzuo L, Ping Y, Xiuli W. Bin zuo hua xue cheng fen Yan jiu. *China J Chin Mater Med*. 2005;30:1390-1391.
140. Metwally MA. Chemical constituents of *Artemisia inculta*. *Boll Chim Farm*. 2001;140(4):265-266.
141. Yinjuan B, Yu L, Yanping S, Youhua H. Chemical composition of *Artemisia argyi*. *Chin Pharm J*. 1997;32:462-465.
142. Abu Zarga M, Qauasmeh R, Sabri S, Munsoor M, Abdalla S. Chemical constituents of *Artemisia arborescens* and the effect of the aqueous extract on rat isolated smooth muscle. *Planta Med*. 1995;61(3):242-245.
143. YH H, Li Y. Chemical constituents of *Artemisia roxburgiana* Wall. *Zhongguo Zhong Yao Za Zhi*. 1994;19(164-165):191.
144. Zequn Y, Hua H, Xiaoqing Z, Ren-Xiang T. Study on chemical constituents of *Artemisia argyi*. *Chin Tradit Herbal Drug*. 1993;24:567-569.
145. Yucheng G, Zuozuo T. Chemical constituents of Japanese wormwood (*Artemisia japonica*). *Chin Tradit Herbal Drug*. 1993;24:122-124.
146. Shi Y. Tie GAN zuo hua xue cheng fen Yan jiu. *Chem J Chin Univ*. 1992;13:1258.
147. Koreeda M, Nagaki M, Hayami KEN-I, Matsueda S. Studies on sesquiterpene lactones. XIII. : Chemical constituents of *Artemisia montana*(NAKAI)PAMP. *Yakugaku Zasshi*. 1988;108(5):434-436.
148. Zan K, Chen X-Q, Zhao M-B, Tu P-F, . Sesquiterpenoids from aerial parts of *Artemisia myriantha*. *Zhongguo Zhong Yao Za Zhi*. 2016;41(15):2833-2837.
149. Liang JY, Ting Liu X. Chemical constituents and insecticidal activity of the essential oils extracted from *Artemisia giraldii* and *Artemisia rubripes* against two stored product insects. *Med chem*. 2016;6(8):541-545.
150. Xiaoyang D, Xinrong D. Study on chemical constituents of young leaves of *Artemisia scoparia* in Hunan. *Northwest Plant J*. 2010;30:1259-1263.
151. Abd-Elhady H. Insecticidal activity and chemical composition of essential oil from *Artemisia judaica* L. against *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). *J Plant Prot Res*. 2012;52(3):348-352.
152. Behtari B, Tilaki GAD, Gholami F, Balkhkanlou RB. Comparison of the essential oil constituents of *Artemisia herba-alba* asso in the vegetative and flowering stages. *Agric Sci Dig*. 2011;31:100-105.
153. Adel K, Zied Z, Ahmed Bkir, Neacute ji G, Mohamed D, Radhouane G. Chemical constituents and antioxidant activity of the essential oil from aerial parts of *Artemisia herba-alba* grown in Tunisian semi-arid region. *Afr J Biotechnol*. 2011;10(15):2923-2929.
154. Taherkhani M. Chemical constituents, antimicrobial, cytotoxicity, mutagenic and antimutagenic effects of *Artemisia ciniformis*. *Iran J Pharm Res*. 2016;15(3):471-481.
155. Judzentiene A, Budiene J. Chemical polymorphism of essential oils of *Artemisia vulgaris* growing Wild in Lithuania. *Chem Biodivers*. 2018;15(2):e1700257.
156. Yan Z. Study on the volatile oil constituents of *Artemisia argyi*. *Guangxi Zhinw*. 2006;26:110-112.
157. Alizadeh M, Aghaei M, Sharifian I, Saadatian M. Chemical composition of essential oil of *Artemisia vulgaris* from West Azerbaijan, Iran. *Electr J Environ Agri Food Chem*. 2012;11:493-496.
158. Huang J, Qian C, Xu H, Huang Y. Antibacterial activity of *Artemisia asiatica* essential oil against some common respiratory infection causing bacterial strains and its mechanism of action in *Haemophilus influenzae*. *Microb Pathog*. 2018;114:470-475.
159. Liang J-Y, Wang W-T, Zheng Y-F, et al. Bioactivities and chemical constituents of essential oil extracted from *Artemisia anethoides* against two stored product insects. *J Oleo Sci*. 2017;66(1):71-76.
160. Liang J-Y, Guo S-S, Zhang W-J, et al. Fumigant and repellent activities of essential oil extracted from *Artemisia dubia* and its main compounds against two stored product pests. *Nat Prod Res*. 2017;32(10):1234-1238.
161. Taherkhani M. Chemical constituents, total phenolic content, antimicrobial, antioxidant and radical scavenging properties, chelating ability, tyrosinase inhibition and in vitro cytotoxic effects of *Artemisia Aucheri* herbs. *Pharm Chem J*. 2017;50(11):736-745.
162. Boghozian A, Amjad L, Shahanipour K. Chemical constituents and identification of the essential oil of *Artemisia aucheri* Boiss. in Iran. *Adv Environ Biol*. 2014;8:2339-2343.
163. Mishra T, Srivastava M, Kumar A, Pal M, Tewari SK. Chemical composition and termiticidal activity of *Artemisia nilagirica* essential oil growing in southern hilly regions of India. *J Essent Oil Bear Pl*. 2017;20(1):247-252.
164. Akhbari M, Aghajani Z, Esmaeili B. Comparative study of antioxidant activity of extracts and essential oils composition of *Artemisia absinthium* L. prepared by two different methods. *J of Essent Oil Bear Pl*. 2014;17(5):954-959.
165. Jihong F, Lijing Z. GC-MS analysis of the volatile oil of Uyghur medicine. *Chinese Journal of Modern Applied Pharmacy*. 2007;24:493-495.
166. Vidic D, Čavar Zeljković S, Dizdar M, Maksimović M. Essential oil composition and antioxidant activity of four *Asteraceae* species from Bosnia. *J Essent Oil Res*. 2016;28(5):445-457.
167. Zanoosi MBP, Nekoei M, Mohammadhosseini M. Composition of the essential oils and volatile fractions of *Artemisia absinthium* by three different extraction methods: Hydrodistillation, solvent-free microwave extraction and headspace solid-phase

- micro extraction combined with a novel QSRR evaluation. *J Essent Oil-Bear Plants*. 2016;19(7):1561-1581.
168. Vieira TM, Dias HJ, Medeiros TCT, et al. Chemical Composition and Antimicrobial Activity of the Essential Oil of *Artemisia absinthium* Asteraceae Leaves. *J Essent Oil Bear Pl*. 2017;20(1):123-131.
169. Kumlay AM, Yildirim BA, Ekici K, Ercisli S. Screening biological activity of essential oils from *Artemisia dracunculus* L. *Oxid Commun*. 2015;38:1320-1328.
170. Verma MK, Anand R, Chisti AM, et al. Essential oil composition of *Artemisia dracunculus* L. (Tarragon) growing in Kashmir-India. *J Essent Oil Bear Pl*. 2010;13(3):331-335.
171. Zhang Y, Zhang J, Yao J, Yang Y-L, Wang L, Dong L-N. Studies on the chemical constituents of the essential oil of *Artemisia dracunculus*. *Zhongguo Zhong Yao Za Zhi*. 2005;30(8):594-596.
172. Radulović NS, Randjelović PJ, Stojanović NM, et al. Toxic essential oils. Part II: chemical, toxicological, pharmacological and microbiological profiles of *Artemisia annua* L. volatiles. *Food Chem Toxicol*. 2013;58:37-49.
173. Xinjian X, Hai S, Guoqing X, Rugang F, Wenfang W. Analysis of chemical constituents of volatile oil from *Artemisia annua* L. by gas chromatography-mass spectrometry. *Lishizhen Med Mater Med Res*. 2009;20:931-932.
174. Zhannan Y, Zhengwen Y, Shiqiong L, Quancai P. Study on the essential oil composition of *Artemisia annua* L. *Lishizhen Med Mater Med Res*. 2008;19:255-257.
175. Rui-zhen L, Yong W, Wei LH. Study on chemical constituents of volatile oil from wild *Artemisia annua* seeds. *Zhongnan Pharmaceutical*. 2007;5:230-232.
176. Zhang Y, Zhang J, Yao J, Wang L, Huang A, Dong L. Studies on the chemical constituents of the essential oil of *Artemisia annua* L. in Xinjiang. *J Northwest Norm Univ*. 2004;40(67-69):86.
177. Creed C, Mollhagen A, Mollhagen N, Pszczolkowski MA. *Artemisia arborescens* "Powis Castle" extracts and  $\alpha$ -thujone prevent fruit infestation by codling moth neonates. *Pharm Biol*. 2015;53(10):1458-1464.
178. You C, Guo S, Zhang W, et al. Identification of repellent and insecticidal constituents from *Artemisia mongolica* essential oil against *Lasioderma serricorne*. *J Chem*. 2015:1-7.
179. Yang C, Hu D-H, Feng Y. Antibacterial activity and mode of action of the *Artemisia capillaris* essential oil and its constituents against respiratory tract infection-causing pathogens. *Mol Med Rep*. 2015;11(4):2852-2860.
180. Wang YL, Zhu DH, Feng XL, YB H, Huang LF. Analysis of the volatile constituents in the dried shoots of *Artemisia capillaris* Thunb. by GC-MS, smoothing and SFA methods. *Chin J Pharm Anal*. 2013;33:98-102.
181. Taherkhani M. Chemical constituents and in vitro anticancer, cytotoxic, mutagenic and antimutagenic activities of *Artemisia diffusa*. *Pharm Chem J*. 2015;48(11):727-732.
182. Liu XC, Li YP, Li HQ, et al. Identification of repellent and insecticidal constituents of the essential oil of *Artemisia rupestris* L. aerial parts against *Liposcelis bostrychophila* Badonnel. *Molecules*. 2013;18(9):10733-10746.
183. Bicchi C, Frattini C, Sacco T. Essential oils of three asiatic *Artemisia* species. *Phytochemistry*. 1985;24(10):2440-2442.
184. Liang C, Hailiang P, Zhenbo G, et al. Study on identification and antioxidant activity of volatile oil from *Herba Artemisiae Scopariae*. *Ziyuan Kaifa Yu Shichang*. 2013;29:469-471.
185. Singh HP, Kaur S, Mittal S, Batish DR, Kohli RK. Essential oil of *Artemisia scoparia* inhibits plant growth by generating reactive oxygen species and causing oxidative damage. *J Chem Ecol*. 2009;35(2):154-162.
186. Xiaoyun Y, Qunfang Q, Zhidan C, Binyin Y. Study on chemical constituents and antioxidant activity of volatile oil from *Artemisia scoparia* Maxim. *Food Technol*. 2012;37:213-217.
187. Zhigzhitzhapova SV, Randalova TE, Radnaeva LD. Composition of essential oil of *Artemisia scoparia* Waldst. et Kit. from Buryatia and Mongolia. *Russ J Bioorgan Chem*. 2016;42(7):730-734.
188. Rana VS, Juyal JP, Blazquez MA, Bodakhe SH. Essential oil composition of *Artemisia parviflora* aerial parts. *Flavour Fragr J*. 2003;18(4):342-344.
189. Shah GC, Bhandari NS, Dhyani P. Essential oil constituents of *Artemisia parviflora* Roxb. *Indian Perfumer*. 2012;56:37-39.
190. Tewari K, Tewari G, Pande C, Kunwar G. Volatile constituents of *Artemisia parviflora* Buch.-Ham. ex Roxb. from Kumaun Himalayan Region, India. *J Essent Oil Bear Pl*. 2015;18(1):195-198.
191. Shah GC, Mathela CS. Investigation on Himalayan *Artemisia* Species VI: Essential Oil Constituents of *Artemisia myriantha* Wall. ex Bess. var. *pleiocephala* (Pamp.) Ling. *J Essent Oil Bear Pl*. 2006;18(6):633-634.
192. Huang H-C, Wang H-F, Yih K-H, Chang L-Z, Chang T-M. Dual bioactivities of essential oil extracted from the leaves of *Artemisia argyi* as an antimelanogenic versus antioxidant agent and chemical composition analysis by GC/MS. *Int J Mol Sci*. 2012;13(11):14679-14697.
193. Meibing L, Yongli Y, Wei L, Maosheng L, Yanfang Z, Huiying Y. Study on chemical constituents and genetic toxicity of volatile oil from leaves of *Artemisia argyi*. *Cbi J Exper Form*. 2012;18:252-255.
194. Xiaowei G, Zongyou L, Peng G, Wei X. Analysis of volatile oil constituents from wild leaves of Jiangsu. *J Beijing Uni Univ*. 2010(24):35-39.
195. Xinjian X, Hai S, Yuqi H, Wenlong Y, Li Z. Analysis of chemical constituents of volatile oil from *Artemisia argyi* with gas chromatography-mass spectrometry. *Lishizhen Med Mater Med Res*. 2007;18:2657-2658.
196. Khan M, Mousa AA, Syamasundar KV, Alkathlan HZ. Determination of chemical constituents of leaf and stem essential oils of *Artemisia monosperma* from central Saudi Arabia. *Nat Prod Commun*. 2012;7(8):1079-1082.
197. Yousefzadeh N. Quantitative and qualitative study of bioactive compounds of essential oils of the medicinal plant *Artemisia sieberi* grown in Lorestan (Iran) by use of GC-MS technique. *Org Chem: Curr Res*. 2012;1(4):109.
198. Ghasemi E, Yamini Y, Bahramifar N, Sefidkon F. Comparative analysis of the oil and supercritical CO<sub>2</sub> extract of *Artemisia sieberi*. *J Food Eng*. 2007;79(1):306-311.



199. Yoon WJ, Moon JY, Song G, et al. *Artemisia fukudo* essential oil attenuates LPS-induced inflammation by suppressing NF- $\kappa$ B and MAPK activation in RAW 264.7 macrophages. *Food Chem Toxicol.* 2010;48(5):1222-1229.
200. Korolyuk EA, Tkachev AV. Chemical composition of the essential oil from two wormwood species *Artemisia frigida* and *Artemisia argyrophylla*. *Russ J Bioorgan Chem.* 2010;36(7):884-893.
201. Qingyuan Z, Shunying Z, Yang Y, Yan Y, Guolin Z. Chemical composition and antimicrobial activity of the essential oil of *Artemisia imponens*. *Journal of Wuban University.* 2009;55:591-596.
202. Chao L, Mingzhu Q. Study on extraction of volatile chemical components from *Artemisia selengensis* by supercritical CO<sub>2</sub> fluid extraction and steam distillation. *Northwest Pharmaceutical Journal.* 2009;24:12-15.
203. Xu Z, Wu Y, Liu D, Chen B. Study on chemical constituents of the essential oil from *Artemisia selengensis* in Dongting lake area. *Se Pu.* 2007;25(5):778-780.
204. Shah GC, Rawat TS. Chemical constituents of *Artemisia indica* Willd. oil. *Indian Perfumer.* 2008;52:27-29.
205. Guibo J, Rensen Z, Shaoxiong C. Identification and antimicrobial effects of volatiles in traditional Chinese medicine herb *Artemisia lavandulaefolia* DC. *Prodr. J Shenyang Agric Univ.* 2008;39:495-498.
206. Ruitao Y, Xiaohui Z, Yanzhen T, Yong S, Lijuan M. Study on chemical constituents of volatile oil from *Artemisia*. *Nat Prod Res Dev.* 2007;19:1005-1008.
207. Ruikar AD, Jadhav RB, Phalgune UD, Rojatkar SR, Puranik VG, Deshpande NR. Phytochemical investigation of *Artemisia pallens*. *Helv Chim Acta.* 2011;94(1):73-77.
208. Changxin A, Weixin Y, Jinjie Z, Niansheng D, Suiguang X. Chemical constituents of essential oil of *Artemisia glauca*. *Chin Tradit Herb Drugs.* 2001;32:591.
209. Lianchang L, Jinan X, Zongcai L, Xiuzhen G. Study on the volatile oil constituents of *Artemisia argyi*. *J Henan Agric Univ.* 1998;32:196-198.
210. Zhixian S, Jinxia Z, Wenhua G, Xizhao Y. Study on chemical constituents of Mongolian *Artemisia* essential oil II. *J Integr Plant Biol.* 1987;29:674-676.
211. Zhixian S, Jinxia Z, Wenhua G, Xizhao Y. Studies on the chemical constituents of Mongolian essential oil I. *Acta Chim Sin.* 1983;41:734-738.
212. Zhou WJ, Zhang SY, Yang YY. Analysis of chemical components of volatile oil from *Artemisia lactiflora* Wall in north Guizhou Province of China. *Med Plant.* 2011;2:59-61.
213. Zhi-Xian S, Xi-Zhao Y. Studies on chemical constituents of the essential oil of *Artemisia subdigitata* Mattf by glass capillary Gas chromatography. *Acta Bot Sin.* 1982;24:159-163.